SIVIOBIO® Small Bio, Smart Tool



CATALOG

& Technical Reference









- · Cloning Vectors
- · DNA Ladders and Dyes
- · Precast Gel
- Protein Markers and Stain
- DNA Amplification
- · cDNA Synthesis and RT-PCR















Original Manufacturing

Table of Contents

Overview of SMOBIO Products

Table of Contents--1

About SMOBIO Technology, Inc.--2

Ordering and General Information--4

Icon Descriptions--5

Tree Diagram of SMOBIO Products--6

Illuminator

Blue Light LED Epi-illuminator--8 (VE0100)

Photobox--9 (VE0104)

Cloning

PCR Cloning Vector--11 (CV1100)

E.coli Transformation Kit--12 (CK1000)

DNA Ladders and Dyes

DNA Ladder--18

(DM1100, DM2100, DM2300, DM3100, DM3200, DM4100, DM5100) (DM1200, DM2000, DM2200, DM2400)

Fluorescent DNA Ladder--24

(DM1160, DM2160, DM2360, DM3160, DM3260, DM4160)

6× DNA Loading Dye--27

(DL1000, DL2000, DL3000, DL4000)

DNA Fluorescent Loading Dye--28

(DL5000, DL5001)

DNA Fluorescent Staining Dye--29

(DS1000, DS1001)

Nucleic Acid Gel Stain-30

(NS1000, NS1001)

Precast Gel

Bis-Tris Precast Gel--36

(QP2110, QP2120, QP2310, QP2320, QP2510, QP2520)

(QP3110, QP3120, QP3310, QP3320, QP3510, QP3520)

TGN Precast Gel--37

(QP4210, QP4220, QP4510, QP4520)

(QP5210, QP5220, QP5510, QP5520)

Running Buffer Powder--38 (BF2000, BF4000)

Protein Markers and Stain

43--All Blue Protein Marker

(PM1500, PM1600, PM1700)

45--3-color Protein Marker and Ladder

(PM2500, PM2510, PM2600, PM2610, PM2700, PM2800)

(PM5000, PM5100, PM5200)

50--Western Marker

(WM1000)

51--Protein Fluorescent Staining Dye

(PS1000, PS1001)

Deoxynucleotide (dNTP)

54--dNTP Single Solution

(CD3000, CD3001)

(CD4000, CD4001)

(CD5000, CD5001)

(CD6000, CD6001)

(CD7000, CD7001)

55--dNTP Mix

(CD1010, CD1011, CD1020, CD1021)

DNA Amplification

60--High Fidelity PCR

(TF1000, TF3000, TK1000)

63--Standard PCR

(TP1000, TP1100/TP1120, TP1200, TP1260)

67--Extraction-free PCR

(TP2000, TP2100)

69--Hot Start PCR

(TP5000)

70--Real-time PCR

(TQ1100, TQ1101, TQ1110, TQ1200, TQ1201, TQ1210, TQ1211)

(TQ2110)

cDNA Synthesis and RT-PCR

75--cDNA Synthesis

(RP1000, RP1300, RP1400)

76--One-Step RT-PCR

(RP1100)

77--One-Step RT-qPCR

(RQ2110, RQ2200)

79--RNase Inhibitors (RI1000, RI1001)

Pathogen Detection

80--COVID-19 Detection

(IP2000)



About SMOBIO Technology, Inc.



Established in 2004, SMOBIO excels in developing and manufacturing molecular biology reagents and testing tools. SMOBIO's motto — "Small Bio, Smart tool" suggests, we take pride in the high quality reagents we make which are specially designed to meet the individual needs of the researchers using them.

Our high quality protein markers contain ready-to-use stabilized combinations of pre-stained proteins with a broad range of molecular weights. They are suitable for just about all protein electrophoresis applications.

In addition, our user-friendly and highly consistent DNA molecular weight markers include a wide selection of DNA ladders ranging from 50 bp to 25 kbp in various packages. SMOBIO also provides DNA polymerase and reverse transcriptase in various package formats. They can be optimized for different laboratory applications, facilitating easy and routine performance of successful PCR, and RT-PCR. Furthermore, we provide safe and highly sensitive DNA and protein fluorescent dyes that can be illuminated by safer LED blue light instead of damaging UV light, resulting in an unprecedented level of safety for the user as well as the highest level of detection sensitivity at the same time.

In an effort to protect users and their samples, SMOBIO created the bi-functional B-BOX™ Blue Light LED epi-illuminator which works flawlessly with DNA and protein related products. B-BOX™ features a multi-angled filter plate that can be adjusted to create a very accessible working area, allowing for real time visualization, precision gel cutting, and documentation all in one compact lightweight unit.





B-BOX™ works without limits using many of SMOBIO's products including SMOBIO's ExcelTaq™ Fluorescent PCR Master Mix, FluoroDye™ DNA Fluorescent Loading Dye, or even the FluoroStain™ DNA Fluorescent Staining Dye for maximum visualization.

DNA in gels with fluorostain will become a better prospect for cloning when used with B-BOX™. Unlike traditional UV light which results in damage to samples, the blue light system allows for safer illumination that doesn't damage biomolecules.

If you are looking for unsurpassed cloning efficiency just pair our B-BOX™ with SMOBIO's GetClone™ PCR Cloning Vector and your experiments will thank you.

Ordering and General Information

Contacting SMOBIO Technology, Inc.

For orders, quotations, or information, please visit www.smobio.com or contact your authorized SMOBIO distributor.

Distributors

For information on finding your nearest SMOBIO distributors, please visit www.smobio.com/distributor.html or email us: info@smobio.com

How to Place an Order or Request a Quote

Quotation requests can be submitted online at www.smobio.com from anywhere in the world or through your authorized SMOBIO distributor.

Prices and Terms

For current prices, contact your authorized SMOBIO distributor.

Perishable and Hazardous Materials Surcharges:

Orders for products classified as dangerous, and those requiring special packaging with blue ice (wet ice) or dry ice may incur additional charges. Please contact your authorized SMOBIO distributor for more information.

Customer Service & Technical Support

For troubleshooting or further information about products, techniques, or applications please visit our **website**:

www.smobio.com

- Product Information, Instruction Manuals, and Brochures
- Technical Data Sheets (TDS)
- Safety Data Sheet(SDS)
- Frequently Asked Questions (FAQs)
- Contact Information





Ordering and General Information





SMOBIO Technology, Inc. will ship orders promptly. If any error or damage occurs, please contact the SMOBIO office immediately (info@smobio.com). Report all shipping damage to the courier, take pictures, and keep all containers and packing material until an inspection is made. If any merchandise must be returned, please contact SMOBIO (info@smobio.com) for instruction.

SMOBIO will not accept returns without prior authorization. Merchandise is returnable for replacement only if requested within 7 days of receiving products, and only if the merchandise remains unused, unopened, and inside the original packaging. SMOBIO cannot authorize any return of products for any erroneously placed order from a distributor.

Please provide the following information for returning merchandise.



Warranty Information

 $\ensuremath{\mathsf{AII}}$ SMOBIO products are guaranteed to meet the specifications listed.

Instruments generally receive 1 year of warranty coverage from the date of shipment or installation. Warranties do not cover consumable parts, such as lamps or platinum wire. Repairs have different warranty periods. Warranties are not transferable from the original purchaser.

SMOBIO is responsible for the quality of its products. We will replace any merchandise with any defect due to a manufacturing error or inadequate precaution made during packaging by sending an additional replacement. The warranty period for B-BOX™ Blue Light LED epi-illuminator is one year.



Icon Descriptions



Tree Diagram of SMOBIO Products







Apparatus

Blue Light LED Epi-illuminator Phox



Cloning

PCR Cloning E.coli Transformation Kit Getclone



DNA **DNA Ladder**

Fluorescent DNA Ladder 6× DNA Loading Dye **DNA Fluorescent Loading** DNA Fluorescent Staining Dye **Nucleic Acid Gel Stain**



Precast Gel

Bis-Tris Precast Gel Tris-Glycine Precast Gel **Running Buffer Powder**



Protein

All Blue Protein Marker 3-color Protein Marker/Ladder

Western Marker **Protein Fluorescent** Staining Dye



dNTP

dNTP Single Solution dNTP Mix



PCR

High Fidelity PCR Standard PCR **Extraction-free PCR Hot Start PCR** Real-time PCR



RNA

cDNA Synthesis One-Step RT-PCR One-Step RT-qPCR **RNase Inhibitors**



Pathogen Detection COVID-19 Detection

Cat. No.IP2000



Blue Light LED Epi-illuminator

Photobox



B-BOX™Blue Light LED Epi-illuminator

VE0100

Description

SMOBIO'S B-BOX™ is a long wavelength, blue light LED epi-illuminator. It is compact in design and robust in construction. The B-BOX™ epi-illuminator provides an unprecedented level of safety for its user due to its non-UV light source and a low operating voltage of only 12 Volts, as well as its capability in working with non-carcinogenic DNA/ protein dye. The B-BOX™ comes with high quality non-flickering LED light that is highly sensitive for DNA and protein dye. B-BOX™ can greatly ease the routine chore of gel extraction, enabling easy visualization and gel cutting even in bright ambient light. The multi-angle filter plate (Cat. No. VE0102) provides optimal angles for gel cutting, visualization, and documentation. The B-BOX™ also comes with amber colored filter goggles (Cat. No. VE0103) for gel cutting and visualization. The Phox™ Photobox (Cat. No. VE0104) provides a mini darkroom environment for images taken with any smartphone. Finally, a built-in barrier system around the working area helps facilitate cleaning.

Features

- Improved cloning efficiency
- Compact, lightweight, and portable (less than 1 kg (in weight)
- Safety features include 470 nm long wavelength, without any UV radiation hazard to its user
- Compatible with non-carcinogenic, non-ethidium bromide DNA staining dye
- User friendly: Samples are easy to visualize (when using the filter plate or goggles)
- LED light source lasts up to 50,000 hours
- Superior detection sensitivity: ≤ 0.04 ng of DNA when using FluoroStain™ DNA Fluorescent Staining Dye, ≤3 ng of
 protein when using FluoroStain™ Protein Fluorescent Staining Dye (as sensitive as silver stain)
- · Adjustable and removable filter plate allows for gel cutting, visualization, and documentation
- Built-in barrier design, for easy clean up
- Visible in bright ambient light
- Emphasizes minimal power reliance, low heat generation, with its own built-in heat sink

Physical Specifications

• Overall Dimensions (mm): 201.4 x 200 x 38 (D x W x H)

• Viewing Area (mm): 158 x 96 (D x W)

• Wavelength of LEDs (nm): 470

Number of LED Units: 72 Super Flux LEDs
 LED Life up to 50,000 hours
 Power: 12 Volt DC, 0.72 Amp

• Electrical Requirements: AC 100~240 V, 50/ 60 Hz (Adapter)

• Weight (kg): 0.95kg (Net Weight)

• Shipping Weight (kg): 1.0 (Gross Weight) – Adapter (0.5 kg) not included

Material: ASA for housing; Tempered glass working area


• Recommended Dyes: ExcelDye™ DNA Fluorescent Loading Dye

FluoroDye™ DNA Fluorescent Staining Dye

FluoroStain™ Protein Fluorescent Staining Dye

FluoroVue™ Nucleic Acid Gel Stain

SYBR Green I Nucleic Acid Gel Stain







Phox™ PhotoBox

VE0104

Description

SMOBIO's Phox[™] is a lightweight portable photobox designed to help users take high quality photos of gels observed with SMOBIO's B-BOX[™] Blue light epi-illuminator. When working with the Phox[™] Photobox, users don't need to turn off any lights or take the gel and system into a dark room to prevent light interference. With a large platform, any smartphone can easily fit the Phox[™] Photobox. the Phox[™] Photobox is designed to fit B-Box[™], but its compact size allows it to work with other UV & blue light illuminators without the need for a large and expensive imaging system.

Features

- 1. Light and portable
- 2. Compatible with B-BOX[™] and various illuminators
- 3. Designed with 3D support structure
- 4. Large platform designed for bigger smartphones
- 5. Made from a single sheet of paper

Physical Specifications

• Weight: 34 g (net weight)

• Size (mm): 155 x145 x 100 (D x W x H)

• Tolerance Weight : 1 kg
• Material : Paper









GetClone™ PCR Cloning Vector II

CV1100 (20 RXN)


Description

The GetClone™ PCR Cloning Vector is a positive selection system for high efficiency cloning of blunt end DNA or amplicons. This cloning vector contains a lethal gene which can be disrupted by ligation of a blunt end DNA insert at the cloning site. Only colonies with inserted vectors are able to propagate, eliminating the need for IPTG and X-Gal for blue/white screening. The GetClone™ pGet II vector includes ampicillin and kanamycin resistance genes that can meet the needs of most users.

Features

- 1. Cloning efficiency greater than 90%
- 2. IPTG and X-Gal are not required
- 3. Accepts a wide range of insert/vector ratios 0.5:1 to 12:1
- 4. Accepts insert sizes of 6 bp to 12 kb
- 5. The phosphorylation of PCR products is not required
- 6. Accepts blunt end amplicon or DNA fragment (not for sticky ends)
- 7. Resistance to ampicillin and kanamycin

Cloning sites of GetClone™ PCR Cloning Vector



CV1100

GetClone™ PCR Cloning Vector II

Contents

| Component | Volume |
|-------------------------|--------|
| pGet II Vector | 23 μΙ |
| pGet-For Primer (10 μM) | 100 µl |
| pGet-Rev Primer (10 μM) | 100 µl |
| | |

Storage -20°C

for 24 months




Fig. 1. Cloning map of pGet II vector.

The pGet II vector carries a pUC origin, two antibiotic resistant genes as selectable markers, and a lethal gene disrupted by cloning sites.





Easy . Fast . Efficient

- Flexible fresh culture, overnight culture, 4°C stored liquid culture or even colonieson agar plate can be used for transformation.
- Fast and Easy only few steps for preparation; suitable for time-saving transformation
- High efficiency up to 10° cfu/μg
- Personalization suitable for most E. coli strains

Description

Champion™ E. coli Transformation Kit provides an easy method for rapid preparation of chemically competent cells with high transformation efficiency from fresh culture, overnight culture, or even directly from bacterial colonies on the plate. The competent cell preparation method eliminates the requirement of time-wasting wash step. In addition, preparation of competent cells from overnight culture or directly from bacterial colonies provides flexibility to cloning experiments. The resultant competent cells can be immediately used or stored at -70°C for one year.

This kit includes a specialized SMO-Broth™ medium and a unique Champion™ CC Buffer for culturing and preparing competent cells efficiently. Following the simple and quick competent cell preparation protocol from fresh culture, the transformation efficiency is typically ranged from 108 to 109 transformants/µg of pUC19 plasmid DNA, but varies depending on the *E. coli* strains.

The resultant competent cells can be further transformed using time-saving transformation protocol, eliminating the requirement of heat-shock and recovery steps.



So Easy to Have
Your Own Competent Cells

Kit Contents

| Component | Volume |
|--|------------|
| Champion™ CC Buffer | 20 ml |
| SMO-Broth™ | 100 ml x 2 |
| pUC19 Control Plasmid (10 ⁻⁴ μg/μl) | 5 μΙ |
| Instruction Manual | 1 |
| Champion™ Competent Cell Preparation Car | rd 1 |

Storage

for 12 months






DNA Ladders and Dyes


DNA Ladder


Fluorescent DNA Ladder 6× DNA Loading Dye DNA Fluorescent Loading Dye DNA Fluorescent Staining Dye Nucleic Acid Gel Stain




DNA Ladders and Markers Information











Fluorescent DNA Ladders







DNA loading dye Information


| Cat.No. | Application | Dilution fold | Suitable gel | Dye component | Dye migration (1% agarose gel/ 1X TAE) |
|---------|-------------|------------------|------------------------------------|--|---|
| DL1000 | DNA | 6X | agarose gel, polyacrylamide gel | Orange G | ~50 bp |
| DL2000 | DNA | 6X | agarose gel, polyacrylamide gel | Orange G, Xylene cyanol FF | ~50 bp, ~4000 bp |
| DL3000 | DNA | 6X | agarose gel, polyacrylamide gel | Bromophenol blue, Xylene cyanol FF | ~500 bp, ~4000 bp |
| DL4000 | DNA | 6X | agarose gel, polyacrylamide gel | Orange G, Bromophenol blue, Xylene cyanol FF | ~50 bp, ~500 bp, ~4000 bp |

DNA fluorescent loading and staining dye Information

| Cat.No. | Application | Dilution fold | Sensitivity | Excitation (nm) | Emission (nm) | Observation |
|------------------|-------------|------------------|-------------|-----------------|---------------|--------------|
| DL5000 DL5001 | DNA | 6X | 0.14 ng | 270, 370, 479 | 522 | B-BOX/UV-BOX |
| DS1000 DS1001 | DNA | 10000X | 0.04 ng | 270, 370, 479 | 522 | B-BOX/UV-BOX |
| NS1000 NS1001 | DNA/RNA | 10000X | 0.14 ng | 250, 482 | 509 | B-BOX/UV-BOX |



Post stain



Premix DNA sample with DL5000

How to use DNA fluorescent loading and staining dye



Running buffer stain



In-gel stain





ExcelBand™ 50 bp DNA Ladder










DM1100 (500 µl)



Description

The DM1100 ExcelBand™ 50 bp DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with loading dye for direct gel loading. The DM1100 DNA ladder is composed of 17 individual DNA fragments: 1.5k, 1.2k, 1k, 900, 800, 700, 600, 500, 450, 400, 350, 300, 250, 200, 150, 100 and 50 bp derived from a mixture of PCR products and specifically digested plasmid DNA. This product contains two enhanced bands (500 bp and 200 bp) for easy reference. In addition, the low range Orange G tracking dye mimics the migration of a 50 bp dsDNA during electrophoresis, and allows for real time monitoring.

Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA.

Range

50 ~ 1,500 bp

Concentration

54 μg/500 μl

Recommended loading volume


5 μl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 36 months

ExcelBand™ 100 bp DNA Ladder





Description

The DM2100 ExcelBand™ 100 bp DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with loading dye for direct gel loading. The DM2100 DNA marker is composed of 11 individual DNA fragments: 1.5k, 1k, 900, 800, 700, 600, 500, 400, 300, 200 and 100 bp derived from a mixture of PCR products and specifically digested plasmid DNA. This product contains two enhanced bands (1.5 kb and 500 bp) for easy reference. In addition, two tracking dyes Xylene cyanol FF and Orange G which mimic the migration of 4,000 bp and 50 bp dsDNA during electrophoresis. These dyes allow for real time monitoring.









Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA

Range

100 ~ 1,500 bp

Concentration

52.2 μg/500 μl

Recommended loading volume 5 µl/well

Storage

Room temperature for 6 months

4°C for 12 months

-20°C for 36 months







ExcelBand™ 100 bp+3K DNA Ladder










DM2300 (500 µl)



Description

The DM2300 ExcelBand™ 100 bp+3K DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with loading dye for direct gel loading. The DM2300 DNA ladder is composed of 12 individual DNA fragments: 3k, 1.5k, 1k, 900, 800, 700, 600, 500, 400, 300, 200 and 100 bp derived from a mixture of PCR products and specifically digested plasmid DNA. This product contains two enhanced bands (1.5 kb and 500 bp) for easy reference. In addition, two tracking dyes, Xylene cyanol FF and Orange G which mimic the migration of 4,000 bp and 50 bp dsDNA during electrophoresis are added for real time monitoring.

Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA.

Range

100 ~ 3,000 bp

Concentration

56 μg/500 μl

Recommended loading volume


5 μl/well

Storage

Room temperature for 6 months for 12 months -20°C for 36 months

ExcelBand™ 1 KB (0.25-10 kb) DNA Ladder





Description

The DM3100 ExcelBand™ 1 KB (0.25-10 kb) DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with loading dye for direct gel loading. The DNA ladder DM3100 is composed of 13 individual DNA fragments: 10k, 8k, 6k, 5k, 4k, 3k, 2.5k, 2k, 1.5k, 1k, 750, 500, and 250 bp derived from a mixture of PCR products and specifically digested plasmid DNA. This product contains two enhanced bands (3 kb and 1 kb) for easy reference. In addition, two tracking dyes, Xylene cyanol FF and Bromophenol blue which mimic the migration of 4,000 bp and 500 bp dsDNA during electrophoresis are added for real time monitoring.









Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA.

Range

250 ~ 10,000 bp

Concentration

50 μg/500 μl

Recommended loading volume

5 μl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 36 months





ExcelBand™ 1 KB Plus (0.1-10 kb) DNA Ladder


DM3200 (500 µI)











Description

The DM3200 ExcelBand™ 1 KB Plus (0.1-10 kb) DNA Ladder is a ready-touse DNA ladder, which is pre-mixed with loading dye for direct gel loading. The DNA Ladder DM3200 is composed of 19 individual DNA fragments: 10k, 8k, 6k, 5k, 4k, 3k, 2.5k, 2k, 1.5k, 1k, 900, 800, 700, 600, 500, 400, 300, 200, and 100 bp derived from a mixture of PCR products and specifically digested plasmid DNA. This product contains three enhanced bands (3 kb, 1.5 kb and 500 bp) for easy reference. In addition, three tracking dyes, Xylene cyanol FF, Bromophenol blue and Orange G which mimic the migration of 4,000 bp, 500 bp and 50 bp dsDNA during electrophoresis are added for real time monitoring.

Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA.

Range

100 ~ 10,000 bp

Concentration

87 μg/500 μl


Recommended loading volume 5 µl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 36 months

ExcelBand™ XL 25 kb DNA Ladder, Broad Range (up to 25 kb)

DM4100 (500 μl)



Description

The DM4100 ExcelBand™ XL 25 kb DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with loading dye for direct gel loading. The DNA Ladder DM4100 is composed of 14 individual DNA fragments: 25k, 10k, 8k, 6k, 5k, 4k, 3k, 2.5k, 2k, 1.5k, 1k, 750, 500, and 250 bp derived from a mixture of PCR products and specifically digested plasmid DNA. This product contains two enhanced bands (3 kb and 1 kb) for easy reference. In addition, two tracking dyes, Xylene cyanol FF and Bromophenol blue which mimic the migration of 4,000 bp and 500 bp dsDNA during electrophoresis are added for real time monitoring.









Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA.

Range

250 ~ 25,000 bp

Concentration

51.6 μg/500 μl


Recommended loading volume 5 µl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 36 months







ExcelBand™ Super Range DNA Ladder (50 bp-25 kb)

DM5100 (500 µl)









Description

The DM5100 ExcelBand™ Super Range DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with loading dye for direct gel loading. The DNA Ladder DM5100 is composed of 26 individual DNA fragments: 25k, 10k, 8k, 6k, 5k, 4k, 3k, 2.5k, 2k, 1.5k, 1.2k, 1k, 900, 800, 700, 600, 500, 450, 400, 350, 300, 250, 200, 150, 100 and 50 base pairs derived from a mixture of PCR products and specifically digested plasmid DNA. This product contains four enhanced bands (3k. 1.2k, 500 and 200 bp) for easy reference. In addition, three tracking dyes, Xylene cyanol FF, Bromophenol blue and Orange G which mimic the migration of 4,000 bp, 500 bp and 50 bp dsDNA during electrophoresis are added for real time monitoring.

Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA

Range

50 ~ 25,000 bp

Concentration

88.7 μg/500 μΙ

Recommended loading volume

5 μl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 36 months





AccuBand™ 50 bp DNA Ladder II


DM1200 (500 μl)











Description

AccuBand™ 50 bp DNA Ladder II is composed of 13 individual DNA fragments, presenting 1k, 900, 800, 700, 600, 500, 400, 300, 250, 200, 150, 100 and 50bp sharp bands respectively. This product contains 2 enhanced bands (500 bp and 250 bp) for easy band identification. AccuBand™ 50 bp DNA Ladder II is ready-to-use, containing loading buffer with a tracking dye (Orange G). AccuBand™ 50 bp DNA Ladder II provides a sufficient amount of DNA for clear observation of all DNA bands ranging from 50 bp to 1 kb, either in agarose gel or in polyacrylamide gel electrophoresis.

Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCI (pH 8.0) and 10 mM EDTA

Range 50 ~ 1000 bp

Concentration 55.5 μg/500 μl

Recommended loading volume


5 μl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 36 months

AccuBand™ 100 bp DNA Marker II

DM2000 (500 ul)



Description

AccuBand™ 100 bp DNA Marker II is composed of 6 individual DNA fragments, presenting 2k, 1k, 750, 500, 250 and 100 bp sharp bands respectively. This product contains 1 enhanced band (750 bp) for easy band identification. AccuBand™ 100 bp DNA Marker II is ready-to-use, containing loading buffer with dual color tracking dyes (Orange G and Xylene cyanol FF). To improve the faint visibility of low molecuweight bands frequently occurred in use of conventional DNA Markers, AccuBand™ 100 bp DNA marker II provides a sufficient amount of DNA for 250 and 100 bp fragments, ensuring clear observation of all DNA bands ranging from 100 bp to 2 kb, either in agarose gel or in polyacrylamide gel electrophoresis.









Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA

Range 100 ~ 2000 bp

Concentration

45.5 μg/500 μΙ

Recommended loading volume 5 μl/well

Storage

Room temperature for 6 months for 12 months -20°C for 36 months





AccuBand™ 100 bp DNA Ladder II










DM2200 (500 µl)



Description

AccuBand™ 100 bp DNA Ladder II is composed of 10 individual DNA fragments, presenting 1k, 900, 800, 700 600, 500, 400, 300, 200 and 100 bp sharp bands respectively. This product contains 1 enhanced band (500 bp) for easy identification of bands. AccuBand™ 100 bp DNA Ladder II is ready-to-use, containing loading buffer with dual color tracking dye (Orange G and Xylene Cyanol FF). AccuBand™ 100 bp DNA Ladder II provides a sufficient amount of DNA for clear observation of all DNA bands ranging from 100 bp to 1 kb, either in agarose gel or in polyacrylamide gel electrophore-

Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA

Range

100 ~ 1000 bp

Concentration

50 μg/500 μl

Recommended loading volume


5 μl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 36 months

AccuBand™ 100 bp+3K DNA Ladder II

DM2400 (500 ul)



Description

AccuBand™ 100 bp+3K DNA Ladder II is composed of 14 individual DNA fragments, presenting 3k, 2k, 1.5k, 1.2k, 1k, 900, 800, 700 600, 500, 400, 300, 200 and 100 bp sharp bands respectively. This product contains 2 enhanced bands (1k and 500 bp) for easy band identification. AccuBand™ 100 bp+3K DNA Ladder II is ready-to-use, containing loading buffer with dual color tracking dyes (Orange G and xylene Cyanol FF). AccuBand™ 100 bp+3K DNA Ladder II provides a sufficient amount of DNA for clear observation of all DNA bands ranging from 100 bp to 3 kb, either in agarose gel or in polyacrylamide gel electrophoresis.









Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA

Range

100 ~ 3000 bp

Concentration

54.4 μg/500 μl

Recommended loading volume

5 μl/well

Storage

Room temperature for 6 months for 12 months 4°C -20°C for 36 months



FluoroBand™ 50 bp Fluorescent **DNA Ladder**










DM1160 (500 µI)



Description

The DM1160 FluoroBand™ 50 bp Fluorescent DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with high sensitivity DNA binding fluorescent dye and loading dye for direct gel loading. The DNA Ladder DM1160 is composed of 17 individual DNA fragments: 1.5k, 1.2k, 1k, 900, 800, 700, 600, 500, 450, 400, 350, 300, 250, 200, 150, 100, and 50 bp derived from a mixture of PCR products and specifically digested plasmid DNA. These bands can be visualized when illuminated with 470 nm blue or UV light. This product contains two enhanced bands (500 bp and 200 bp) for easy reference. In addition, the low range Orange G tracking dye which mimics the migration of a 50 bp dsDNA during electrophoresis is added for real time monitoring. Real time observation of electrophoresis is also possible if a compatible light source is fitted to the electrophoresis tank.

Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA

Range

50 ~ 1,500 bp

Concentration

54 µg/500 µl


Recommended loading volume 5 μl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 24 months

FluoroBand™ 100 bp Fluorescent **DNA Ladder**

DM2160 (500 µI)



Description

The DM2160 FluoroBand™ 100 bp Fluorescent DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with high sensitivity DNA binding fluorescent dye and loading dye for direct gel loading. The DNA Ladder DM2160 is composed of 11 individual DNA fragments: 1.5k, 1k, 900, 800, 700, 600, 500, 400, 300, 200 and 100 bp derived from a mixture of PCR products and specifically digested plasmid DNA. These bands can be visualized when illuminated with 470 nm blue or UV light. This product contains two enhanced bands (1.5 kb and 500 bp) for easy reference. In addition, two tracking dyes, Xylene cyanol FF and Orange G which mimic the migration of 4,000 bp and 50 bp dsDNA during electrophoresis are added for real time monitoring. Real time observation of electrophoresis is also possible if a compatible light source is fitted to the electrophoresis tank.









Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA.

Range

100 ~ 1,500 bp

Concentration

52.2 μg/500 μl

Recommended loading volume 5 μl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 24 months





FluoroBand™ 100 bp+3K Fluorescent DNA Ladder










DM2360 (500 µl)



Description

The DM2360 FluoroBand™ 100bp+3K Fluorescent DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with high sensitivity DNA binding fluorescent dye and loading dye for direct gel loading. The DNA Ladder DM2360 is composed of 12 individual DNA fragments: 3k, 1.5k, 1k, 900, 800, 700, 600, 500, 400, 300, 200 and 100 bp derived from a mixture of PCR products and specifically digested plasmid DNA. These bands can be visualized when illuminated with 470 nm blue or UV light. This product contains two enhanced bands (1.5 kb and 500 bp) for easy reference. In addition, two tracking dyes, Xylene cyanol FF and Orange G which mimic the migration of 4,000 bp and 50 bp dsDNA during electrophoresis are added for real time monitoring. Real time observation of electrophoresis is also possible if a compatible light source is fitted to the electrophoresis tank.

Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA.

Range

100 ~ 3,000 bp

Concentration

56 μg/500 μl

Recommended loading volume


5 μl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 24 months

FluoroBand™ 1 KB (0.25-10 kb) Fluorescent DNA Ladder

DM3160 (500 µI)



Description

The DM3160 FluoroBand™ 1KB (0.25-10 kb) Fluorescent DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with high sensitivity DNA binding fluorescent dye and loading dye for direct gel loading. The DNA ladder DM3160 is composed of 13 individual DNA fragments: 10k, 8k, 6k, 5k, 4k, 3k, 2.5k, 2k, 1.5k, 1k, 750, 500, and 250 bp derived from a mixture of PCR products and specifically digested plasmid DNA. These bands can be visualized when illuminated with 470 nm blue or UV light. This product contains two enhanced bands (3 kb and 1 kb) for easy reference. In addition, two tracking dyes, Xylene cyanol FF and Bromophenol blue which mimic the migration of 4,000 bp and 500 bp dsDNA during electrophoresis are added for real time monitoring. Real time observation of electrophoresis is also possible if a compatible light source is fitted to the electrophoresis tank.









Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA.

Range

250 ~ 10,000 bp

Concentration

50 μg/500 μl

Recommended loading volume

5 μl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 24 months





FluoroBand™ 1 KB Plus (0.1-10 kb) Fluorescent DNA Ladder










DM3260 (500 µl)



Description

The DM3260 FluoroBand™ 1 KB Plus (0.1-10 kb) Fluorescent DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with high sensitivity DNA binding fluorescent dye and loading dye for direct gel loading. The DNA Ladder DM3260 is composed of 19 individual DNA fragments: 10k, 8k, 6k, 5k, 4k, 3k, 2.5k, 2k, 1.5k, 1k, 900, 800, 700, 600, 500, 400, 300, 200, and 100 bp derived from a mixture of PCR products and specifically digested plasmid DNA; these bands can be visualized when illuminated with 470 nm blue or UV light. This product contains three enhanced bands (3 kb, 1.5 kb and 500 bp) for easy reference. In addition, three dyes, Xylene cyanol Bromophenol blue and Orange G which mimic the migration of 4,000 bp, 500 bp and 50 bp dsDNA during electrophoresis are added for real time monitoring. Real time observation of electrophoresis is also possible if a compatible light source is fitted to the electrophoresis tank.

Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCI (pH 8.0) and 10 mM EDTA.

Range

100 ~ 10.000 bp

Concentration

87 μg/500 μΙ


Recommended loading volume 5 μl/well

Storage

Room temperature for 6 months 4°C for 12 months -20°C for 24 months

FluoroBand™ XL 25 kb Fluorescent DNA Ladder, Broad Range (up to 25 kb)





Description

The DM4160 FluoroBand™ XL 25 kb Fluorescent DNA Ladder is a ready-to-use DNA ladder, which is pre-mixed with high sensitivity DNA binding fluorescent dye and loading dye for direct gel loading. The DNA Ladder DM4160 is composed of 14 individual DNA fragments: 25k, 10k, 8k, 6k, 5k, 4k, 3k, 2.5k, 2k, 1.5k, 1k, 750, 500, and 250 bp derived from a mixture of PCR products and specifically digested plasmid DNA; these bands can be visualized when illuminated with 470 nm blue or UV light. This product contains two enhanced bands (3 kb and 1 kb) for easy reference. In addition, two tracking dyes, Xylene cyanol FF and Bromophenol blue which mimic the migration of 4,000 bp and 500 bp dsDNA during electrophoresis are added for real time monitoring. Real time observation of the electrophoresis is also possible if a compatible light source is fitted to the electrophoresis tank.









Source

Phenol extracted PCR products and dsDNA digested with specific restriction enzymes, equilibrated in 10 mM Tris-HCl (pH 8.0) and 10 mM EDTA

Range

250 ~ 25,000 bp

Concentration

51.6 μg/500 μl

Recommended loading volume 5 μl/well

Storage

Room temperature for 6 months for 12 months -20°C for 24 months








ExcelDye™ 6X DNA Loading Dye


DL1000 (5 ml x2) DL2000 (5 ml x2) DL3000 (5 ml x2) DL4000 (5 ml x2)


Description

The SMOBIO ExcelDye™ 6X DNA Loading Dye is pre-mixed buffer for loading and tracking DNA samples during electrophoresis on agarose or polyacrylamide gels. It contains different composition of three dyes (Xylene cyanol FF, Bromophenol blue, and Orange G) for tracking DNA migration. The Xylene cyanol FF, Bromophenol blue and Orange G migrate at approximately 800 bp, 150 bp and 30 bp on a standard 2% agarose gel/0.5X TAE buffer respectively (4,000 bp, 500 bp and 50 bp on 1% agarose gel/0.5X TAE buffer respectively). The included glycerol keeps the DNA at the bottom of the well and the presence of EDTA chelates divalent metal ions to prevent a process of metal-dependent nuclease.

| DL1000 ExcelDye™ 6X | DNA Load | ing Dye, O | range |
|--|-----------|------------------------|----------|
| Contents | | | |
| Component | | Volu | m e |
| ExcelDye™ 6X DNA Loading Dye | e, Orange | 5 ml > | ζ 2 |
| Composition 0.15% Orange G 10 mM Tris-HCl (pH 8.0) 60% glycerol 60 mM EDTA | bp | | dye |
| Storage 4°C for 12 months -20°C for 36 months | ~50 | | Orange G |
| | 1%Ag | garose Gel,0.5X TAE Bu | iffer |

















DL5000 (1 ml) DL5001 (1 ml × 5)

Description

FluoroDye™ DNA Fluorescent Loading Dye is a ready-to-use 6X DNA loading dye designed for fast qualitative electrophoresis analysis. Containing sensitive fluorescent dye with high specific affinity towards double stranded DNA (dsDNA), the FluoroDye™ DNA Fluorescent Loading Dye has negligible background and renders destaining process unnecessary. The FluoroDye™ DNA Fluorescent Loading Dye allows the user to immediately visualize electrophoresis result upon completion or to monitor the electrophoresis in real time. FluoroDye™ DNA Fluorescent Loading Dye is compatible with both the conventional UV gel-illuminating system as well as the less harmful long wavelength blue light illumination system. FluoroDye™ emission as bound to dsDNA is 522 nm, while its excitation peaks are at 270, 370 and 497 nm (Fig. 1).

Run Jour DNA sample with loading dye only i

Contents

FluoroDye™ DNA Fluorescent Loading Dye is stored in 6X concentration in 60% glycerol and buffered with Tris-HCl and EDTA, containing Bromophenol blue, Xylene cyanol FF, and Orange G as tracking dyes.

Storage

Protected from light
-20°C for 24 months




Fig. 1. DL5000 FluoroDye™ DNA Fluorescent Loading Dye emission as bound to dsDNA is 522 nm while its excitation peaks are at 270, 370 and 497 nm.




Fig. 2. FluoroDye™ DNA Fluorescent Loading Dye is capable of detecting dsDNA fragments down to 0.14 ng.













Description

FluoroStain™ DNA Fluorescent Staining Dye is designed to be a safer replacement for conventional Ethidium bromide (EtBr) which poses a significant health and safety hazard for its users (Fig. 3). FluoroStain™ DNA Fluorescent Staining Dye offers at least 10 times greater sensitivity in DNA detection levels, and is capable of detecting double stranded DNA (dsDNA) fragments up to 0.04 ng in electrophoresis analysis

FluoroStain™ DNA Fluorescent Staining Dye shows a high specificity to dsDNA, with negligible background signal, making the destaining process entirely optional. FluoroStain™ DNA Fluorescent Staining Dye is compatible with both the conventional UV gel-illuminating systems as well as the less harmful long wavelength blue light illumination systems. The emission when bound to dsDNA is 522 nm, while its excitation peaks are at 270, 370 and 497 nm (Fig. 2)

Contents

Proprietary dye in a 10,000X concentration.

Storage

Protected from light 4°C for 12 months -20°C for 24 months




Fig. 1. FluoroStain™ DNA Fluorescent Staining Dye exhibits extreme sensitivity when detecting dsDNA.




Fig. 2. FluoroStain™ DNA fluorescent Staining Dye (Green, 10,000X), emission as bound to dsDNA is 522 nm while its excitation peaks are at 270, 370 and 497 nm.




Fig. 3. FluoroStain™ DNA Fluorescent Staining Dye shows no mutagenic activity in the Ames test.











$NS1000 (500 \mu I) NS1001 (500 \mu I \times 5)$

Description

FluoroVue™ Nucleic Acid Gel Stain (10,000X) is specially designed for in-gel use and is a safer replacement for conventional Ethidium bromide (EtBr), which poses a significant health and safety hazard its users (Fig.1). It is a fluorescent stain which offers high sensitivity detection of double-stranded or single-stranded DNA and RNA in a convenient manner. FluoroVue™ Nucleic Acid Gel Stain offers high sensitivity (Table 1 and Fig.2) that is several times greater than EtBr.

FluoroVue™ Nucleic Acid Gel Stain is compatible with both conventional UV gel-illumination systems as well as harmless long wavelength blue light illumination systems, like B-BOX™. When bound to nucleic acids, FluoroVue™ Nucleic Acid Gel Stain has a fluorescent excitation maximum of ~250 and ~482 nm, and an emission maximum of ~509 nm (Fig. 3). Therefore, it can replace EtBr without the need of changing existing lab imaging systems.




Fig. 1. FluoroVue $^{\rm TM}$ Nucleic Acid Gel Stain shows no mutagenic activity in the Ames test.

Contents

Proprietary dye in a 10,000X concentration.

Storage

Protected from light 4°C for 12 months -20°C for 24 months

Table 1. Different staining methods for using the FluoroVue™ Nucleic Acid Gel Stain

| Staining methods ¹ | Required dye ² | Sensitivity ³ | Convenience |
|---------------------------------|---------------------------|--------------------------|-------------|
| In- gel staining | 4 μΙ | 0.14 ng | Very good |
| Staining during electrophoresis | 30 μΙ | 0.56 ng | Very good |
| Post stain | 10 μΙ | 0.56 ng | Good |

- 1 For detailed protocols of different staining methods: please see the product information. We recommend using an in-gel staining method for optimal effect.
- 2 With a mini horizontal gel electrophoresis system: Combine 40 ml of agarose gel with 300 ml running buffer. The regular post staining buffer volume is 100 ml.
- 3 Sensitivity is evaluated according to the 4 kb band of DM3100.




Fig. 2. The FluoroVue™ Nucleic Acid Gel Stain shows a green-yellow fluorescence under blue light excitation. The sensitivity of NS1000 is about 0.14 ng (arrow) for a 4 kb fragment




Fig. 3. The emission and exitation spectrum of FluoroVue™ Nucleic Acid Gel Stain



















Catalog & Technical Reference



Q-PAGE™ Bis-Tris Precast Gel







(10 gels)

Description

Q-PAGE™ Bis-Tris Precast Gel is a high-performance and easy to use precast polyacrylamide gel for electrophoresis in Bis-Tris buffer system (MOPS or MES). The optimized gel formula allows Q-PAGE™ Bis-Tris Precast Gel to show improved resolution, accurate results, and an extended shelf-life over conventional Laemmli Tris-HCl gels. Q-PAGE™ Bis-Tris Precast Gels are available in gradient (4 to 12%) and fixed (8% and 12%) concentrations of polyacrylamide in 12-and 15-well formats. Two available cassette sizes, Mini (10 x 8.3 cm) and Midi (10 x 10 cm), are compatible with most popular protein electrophoresis systems. Q-PAGE™ Mini (QP2XXX) Gels are suitable for Bio-Rad® and other systems. Q-PAGE™ Midi (QP3XXX) Gels are suitable for Invitrogen® XCell SureLock® Mini-Cell, Invitrogen® Mini Gel Tank, Hoefer SE260, and other systems.

Features

• User-friendly gel cassette:

Numbered and framed wells for sample loading Labeled warning sign and green tape as reminder

• Easy compatibility:

Available as homogeneous and adjusted gradient gels for a wide range of protein separation

Compatible with most popular protein electrophoresis systems

• Enhanced gel performance:

Enhanced band sharpness

Better resolution of small proteins

Stable for shipping at ambient temperature



Product list

| Bis-Tris | | | | | | |
|---------------|---------|---------|---------|---------|--|--|
| Cassette size | Mi | ni | Mi | di | | |
| Well No. | 12 well | 15 well | 12 well | 15 well | | |
| 8% | QP2110 | QP2120 | QP3110 | QP3120 | | |
| 12% | QP2310 | QP2320 | QP3310 | QP3320 | | |
| 4-12% | QP2510 | QP2520 | QP3510 | QP3520 | | |

Storage and stability

Store Q-PAGE™ Precast Gels at 4°C for periods up to 12 months.

Do not freeze Q-PAGE™ Precast Gels. Remove tape and comb before electrophoresis.

※ Keep Q-PAGE™ Precast Gels flat during storage.







Q-PAGE™ TGN Precast Gel







(10 gels)

Description

Q-PAGE™ TGN (Tris-Glycine Novel) Precast Gels are ready-to-use acrylamide gels for SDS-PAGE running in Tris-Glycine buffer system. With unique formula, Q-PAGE™ TGN Precast Gels perform enhanced speed, better separation, and longer shelf life as compared with conventional Laemmli Tris-HCl gels. The protein migration patterns in Q-PAGE™ TGN series, however, are similar with typical Laemmli Tris-HCl gels, and thus Q-PAGE™ TGN Precast Gels are compatible to traditional SDS-PAGE and subsequent analyses.

Q-PAGE™ TGN Precast Gels are available in gradient (4 to 15%) and fixed (10%) concentrations of polyacrylamide in 12and 15-well formats. Two available cassette sizes, Mini (10 x 8.3 cm) and Midi (10 x 10 cm), are compatible with most popular protein electrophoresis systems. Q-PAGE™ Mini (QP4XXX) Gels are suitable for Bio-Rad® and other systems. Q-PAGE™ Midi (QP5XXX) Gels are suitable for Invitrogen® XCell SureLock® Mini-Cell, Invitrogen® Mini Gel Tank, Hoefer SE260, and other systems.

Features

• User-friendly gel cassette:

Numbered and framed wells for sample loading Labeled warning sign and green tape as reminder

Easy compatibility:

Available as homogeneous and adjusted gradient gels for a wide range of protein separation Compatible with most popular protein electrophoresis systems

• Enhanced gel performance:

Enhanced band sharpness Better resolution of small proteins Stable for shipping at ambient temperature



Product list

| | | TGN | | |
|---------------|---------|---------|---------|---------|
| Cassette size | Mi | ini | Mi | di |
| Well No. | 12 well | 15 well | 12 well | 15 well |
| 10% | QP4210 | QP4220 | QP5210 | QP5220 |
| 4-15% | QP4510 | QP4520 | QP5510 | QP5520 |

Storage and stability

Store Q-PAGE™ Precast Gels at 4°C for periods up to 12 months. Do not freeze Q-PAGE™ Precast Gels. Remove tape and comb before electrophoresis.

※ Keep Q-PAGE™ Precast Gels flat during storage.





SMOChem™ 1X MOPS-SDS Running Buffer Powder

BF2000 (For 5000 ml)

Description

1X MOPS-SDS Running Buffer Powder is used for electrophoresis on Q-PAGE™ Bis-Tris Precast Gel or polyacrylamide gels in Bis-Tris buffer system. It is convenient and universal for electrophoresis in Bis-Tris buffer system.

Features

• Reliable:

Rigorous quality control for reproducible separation of protein electrophoresis.

• Convenient:

Premeasured pouches make 1 liter of 1X buffer solution; No pH adjustment is necessary.

Contents

1X MOPS-SDS Running Buffer Powder, 5 packs

• Fast:

Dissolving in minutes and then ready to use.

• Stable:

Powder packaging suitable for long-term storage.

Storage and stability

Room Temperature for 24 months

SMOChem™ 1X Tris-Glycine-SDS Running Buffer Powder

BF4000 (For 5000 ml)

Description

1X Tris-Glycine-SDS Running Buffer Powder is used for electrophoresis on Q-PAGE™ TGN Precast Gel or Laemmli Tris-HCl gels in Tris-Glycine buffer system. It is convenient and universal for electrophoresis in Tris-Glycine buffer system.

Features

• Reliable:

Rigorous quality control for reproducible separation of protein electrophoresis.

• Convenient:

Premeasured pouches make 1 liter of 1X buffer solution; No pH adjustment is necessary.

• Fast:

Dissolving in minutes and then ready to use.

• Stable:

Powder packaging suitable for long-term storage.

Contents

1X Tris-Glycine-SDS Running Buffer Powder, 5 packs

Storage and stability

Room Temperature for 24 months
















Protein Markers and Stain

All Blue Protein Marker
3-color Protein Marker and Ladder
Western Marker
Protein Fluorescent Staining Dye



Overview of Protein Markers and Ladders





Molecular Weight Range of Protein Markers and Ladders

| 5 | 10 | 20 | 30 | 40 | 50 | 60 | 75 | 100 | 140 | 180 | 245 | 310 | kDa |
|---|------------|------|-----|-----|----|----|----|-----|-----|-----|------|---------|-----------|
| | | 2 | 5 | | | | 72 | | | | PM15 | 000 | |
| | | 2 | 5 | | | | 72 | | | | PM16 | 000 | |
| | | 2 | 5 | | | | 72 | | | | | PM: | 1700 |
| | | 2 | 5 | | | | 75 | | | | PM2 | 500/251 | 10 |
| | |) (2 | 25 | | | | 75 | | | | | PM | 2600/2610 |
| | \bigcirc |) (2 | 25) | | | | 75 | | | | | PM | 2700 |
| | |) (2 | 25) | | | | 75 | | | | | 310 | PM2800 |
| | \bigcirc | 20 | | (40 | | | 75 | | | | PM50 | 00 | |
| | \bigcirc | 20 | | (40 | | | 75 | | | | | PN | 15100 |
| | | 20 | | (40 | | | 75 | | | | | PN | 15200 |

Protein Marker and Ladder Information

| Series Name | Catalog Number | MW Range* | Band Number | Band Color | Enhanced (Markered) Bands |
|-------------|----------------|------------|-------------|------------|---------------------------|
| ExcelBand™ | PM1500 | 10~180 kDa | 10 | В | 25, 72 kDa |
| ExcelBand™ | PM1600 | 10~180 kDa | 11 | В | 25, 72 kDa |
| ExcelBand™ | PM1700 | 10~240 kDa | 12 | В | 25, 72 kDa |
| ExcelBand™ | PM2500 | 10~180 kDa | 10 | R/G/B | 25, 75 kDa |
| ExcelBand™ | PM2510 | 10~180 kDa | 10 | R/G/B | 25, 75 kDa |
| ExcelBand™ | PM2600 | 10~245 kDa | 12 | R/G/B | 25, 75 kDa |
| ExcelBand™ | PM2610 | 10~245 kDa | 12 | R/G/B | 25, 75 kDa |
| ExcelBand™ | PM2700 | 5~245 kDa | 13 | R/G/B | 25, 75 kDa |
| ExcelBand™ | PM2800 | 10~310 kDa | 13 | R/G/B | 25, 75, 310 kDa |
| ExcelBand™ | PM5000 | 10~180 kDa | 13 | R/G/B | 40, 75 kDa |
| ExcelBand™ | PM5100 | 10~245 kDa | 14 | R/G/B | 40, 75 kDa |
| ExcelBand™ | PM5200 | 5~245 kDa | 15 | R/G/B | 40, 75 kDa |
| YesBlot™ | WM1000 | 15~200 kDa | 10 | R/G/B | 30, 80 kDa |

^{*}In Tris-Glycine buffer

ExcelBand™ All Blue Regular Range Protein Marker (9-180 kDa)










$PM1500 (250 \mu I \times 2)$



Description

The PM1500 ExcelBand™ All Blue Regular Range Protein Marker is a blue protein standard with 10 pre-stained proteins covering a wide range of molecular weights from 10 to 180 kDa in Tris-Glycine buffer (9 to 170 kDa in Bis-Tris (MOPS) buffer and Bis-Tris (MES) buffer). Proteins are covalently coupled with a blue chromophore, and two reference bands (at 25 kDa and 72 kDa, respectively) are enhanced in intensity when separated on SDS-PAGE (Tris-Glycine buffer).

The PM1500 ExcelBand™ All Blue Regular Range Protein Marker is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.

Contents

Approximately 0.1~0.5 mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5 at 25°C), 2% SDS, 0.2 mM Dithiothreitol, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control

Under suggested conditions, the PM1500 ExcelBand™ All Blue Regular Range Protein Marker resolves 10 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.

Storage


4°C -20°C

for 3 months

for long term storage

ExcelBand™ All Blue Regular Range Plus Protein Marker (9-180 kDa)

$PM1600 (250 \mu I \times 2)$



Tris-Glycine

Description

The PM1600 ExcelBand™ All Blue Regular Range Plus Protein Marker is a blue protein standard with 11 pre-stained proteins covering a wide range of molecular weights from 10 to 180 kDa in Tris-Glycine buffer (9 to 170 kDa in Bis-Tris (MOPS) buffer and Bis-Tris (MES) buffer). Proteins are covalently coupled with a blue chromophore, and two reference bands (at 25 kDa and 72 kDa, respectively) are enhanced in intensity when separated on SDS-PAGE (Tris-Glycine

The PM1600 ExcelBand™ All Blue Regular Range Plus Protein Marker is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.









Contents

Approximately 0.1~0.5 mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5 at 25°C), 2% SDS, 0.2 mM Dithiothreitol, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control

Under suggested conditions, the PM1600 ExcelBand™ All Blue Regular Range Plus Protein Marker resolves 11 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.

Storage

4°C

for 3 months

-20°C

for long term storage





ExcelBand™ All Blue Broad Range Protein Marker (9-240 kDa)










PM1700 (250 µl × 2)



Tris-Glycine

Description

The PM1700 ExcelBand™ All Blue Broad Range Protein Marker is a blue protein standard with 12 pre-stained proteins covering a wide range of molecular weights from 10 to 240 kDa in Tris-Glycine buffer (9 to 235 kDa in Bis-Tris (MOPS) buffer and Bis-Tris (MES) buffer). Proteins are covalently coupled with a blue chromophore, and two reference bands (at 25 kDa and 72 kDa, respectively) are enhanced in intensity when separated on SDS-PAGE (Tris-Glycine buffer).

The PM1700 ExcelBand™ All Blue Broad Range Protein Marker is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.

Contents

Approximately $0.1^{\circ}0.5$ mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5 at 25°C), 2% SDS, 0.2 mM Dithiothreitol, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control

Under suggested conditions, the PM1700 ExcelBand™ All Blue Broad Range Protein Marker resolves 12 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.

Storage

4°C for 3 months -20°C for long term storage



ExcelBand™ 3-color Regular Range Protein Marker (9-180 kDa)










$PM2500 (250 \mu I \times 2)$



Description

The PM2500 ExcelBand™ 3-color Regular Range Protein Marker is a ready-to-use three-color protein standard with 10 pre-stained proteins covering a wide range of molecular weights from 10 to 180 kDa in Tris-Glycine Buffer (9 to 170 kDa in Bis-Tris (MOPS) buffer and 10 to 170 kDa in Bis-Tris (MES) buffer). Proteins are covalently coupled with a blue chromophore except for two reference bands (one green and one red band at 25 kDa and 75 kDa respectively) when separated on SDS-PAGE (Tris-Glycine buffer). PM2500 ExcelBand™ 3-color Regular Range Protein Marker is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.

Contents

Approximately 0.1~0.4 mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5 at 25°C), 2% SDS, 0.2 mM Dithiothreitol, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control

Under suggested conditions, the PM2500 ExcelBand[™] 3-color Regular Range Protein Marker resolves 10 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.


Storage

4°C -20°C for 3 months

for long term storage

ExcelBand™ Enhanced 3-color Regular Range Protein Marker (9-180 kDa)

$PM2510 (250 \mu I \times 2) PM2511 (250 \mu I \times 10)$



Description

PM2510/PM2511 ExcelBand™ Enhanced 3-color Regular Range Protein Marker is a ready to use three-color protein standard with 10 pre-stained proteins covering a wide range of molecular weights from 10 to 180 kDa (9 to 170 kDa in Bis-Tris (MOPS) buffer and 10 to 170 kDa in Bis-Tris (MES) buffer). Proteins are covalently coupled with a blue chromophore except for two reference green and red bands (25 kDa and 75 kDa, respectively) when separated on SDS-PAGE (Tris-Glycine buffer). The PM2510/PM2511 ExcelBand™ Enhanced 3-color Regular Range Protein Marker is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.









Contents

Approximately 0.2~0.6 mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5), 2% SDS, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control

Under suggested conditions, the PM2510/PM2511 ExcelBand™ Enhanced 3-color Regular Range Protein Marker resolves 10 major bands in SDS-PAGE (Tris- Glycine, MOPS, and MES buffer) and after Western blotting transfer to a nitrocellulose membrane.

Storage

4°C

for 3 months

-20°C

for long term storage





ExcelBand™ 3-color High Range Protein Marker (9-245 kDa)










PM2600 (250 µl × 2)



Description

The PM2600 ExcelBand™ 3-color High Range Protein Marker is a ready-to-use three-color protein standard with 12 pre-stained proteins covering a wide range of molecular weights from 10 to 245 kDa in Tris-Glycine Buffer (9 to 235 kDa in Bis-Tris (MOPS) buffer and 10 to 235 kDa in Bis-Tris (MES) buffer). Proteins are covalently coupled with different chromophores for easy identification of bands, with two reference proteins carrying enhanced intensity corresponding to a green at 25 kDa and red at 75 kDa, respectively, as separated on SDS-PAGE (Tris-Glycine buffer). The PM2600 3-color High Range Protein Marker is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.

Contents

Approximately $0.1^{\circ}0.4$ mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5 at 25°C) 2% SDS, 0.2 mM Dithiothreitol, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control


Under suggested conditions, the PM 2600 ExcelBand™ 3-color High Range Protein Marker resolves 12 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.

Storage

4°C for 3 months
-20°C for long term storage

ExcelBand™ Enhanced 3-color High Range Protein Marker (9-245 kDa)

PM2610 (250 µl × 2) PM2611 (250 µl × 10)



Description

The PM2610/PM2611 ExcelBand™ Enhanced 3-color High Range Protein Marker is a ready-to-use three-color protein standard with 12 pre-stained proteins covering a wide range of molecular weights from 10 to 245 kDa in Tris-Glycine Buffer (9 to 235 kDa in Bis-Tris (MOPS) buffer and 10 to 235 kDa in Bis-Tris (MES) buffer). Proteins are covalently coupled with different chromophores for easy identification of bands, with two reference proteins carrying enhanced intensity corresponding to a green at 25 kDa and red at 75 kDa, respectively, as separated on SDS-PAGE (Tris-Glycine buffer). The Excel-Band™ Enhanced 3-color High Range Protein Marker is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.









Contents

Approximately $0.2^{\circ}0.6$ mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5 at 25°C), 2% SDS, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control

Under suggested conditions, the PM2610/PM2611 ExcelBand™ Enhanced 3-color High Range Protein Marker resolves 12 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.

Storage

4°C for 3 months -20°C for long term storage



ExcelBand™ 3-color Broad Range Protein Marker (3.5-245 kDa)










PM2700 (250 µl × 2)



Description

The PM2700 ExcelBand™ 3-color Broad Range Protein Marker is a ready-to-use three-color protein standard with 13 pre-stained proteins covering a wide range of molecular weights from 5 to 245 kDa in Tris-Glycine buffer (3.5 kDa to 235 kDa in Bis-Tris (MOPS) buffer and Bis-Tris (MES) buffer). Proteins are covalently coupled with a blue chromophore except for two reference bands (one green and one red band at 25 kDa and 75 kDa, respectively) when separated on SDS-PAGE (Tris-Glycine buffer). The PM2700 ExcelBand™ 3-color Broad Range Protein Marker is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.

Contents

Approximately 0.1~0.4 mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5 at 25°C), 2% SDS, 0.2 mM Dithiothreitol, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control

Under suggested conditions, the PM 2700 ExcelBand™ 3-color Broad Range Protein Marker resolves 13 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.


Storage

4°C -20°C for 3 months

for long term storage

ExcelBand™ 3-color Extra Range Protein Marker (10-310 kDa)

$PM2800 (250 \mu I \times 2)$



Tris-Glycine

Description

The PM2800 ExcelBand™ 3-color Extra Range Protein Marker is a ready-to-use three-color protein standard with 13 pre-stained proteins covering an extra range of molecular weights from 10 to 310 kDa in Tris-Glycine Buffer (9 to 290 kDa in Bis-Tris (MOPS) buffer and 10 to 290 kDa in Bis-Tris (MES) buffer). Proteins are covalently coupled with a blue chromophore except for three reference bands (one green and two red bands at 25 kDa and 75, 310 kDa respectively) when separated on SDS-PAGE (Tris-Glycine buffer). The PM2800 ExcelBand™ 3-color Extra Range Protein Marker is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.









Contents

Approximately 0.1~0.4 mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5 at 25°C), 2% SDS, 0.2 mM Dithiothreitol, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control

Under suggested conditions, the PM 2800 ExcelBand™ 3-color Extra Range Protein Marker resolves 13 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.

Storage

4°C -20°C for 3 months

for long term storage

ExcelBand™ 3-color Pre-Stained Protein Ladder, Regular Range (9-180 kDa)









PM5000 (250 μ l × 2)

kDa 180 — 140 — 100 — 75 — 60 — 50 — 40 — 35 — 30 — 25 — 20 — 15 —

Tris-Glycine

Description

The PM5000 ExcelBand™ 3-color Pre-Stained Protein Ladder Regular Range is a ready-to-use three-color protein standard with 13 pre-stained proteins covering a wide range of molecular weights from 10 to 180 kDa in Tris-Glycine Buffer (9 to 170 kDa in Bis-Tris (MOPS) buffer and 10 to 170 kDa Bis-Tris (MES) buffer). Proteins are covalently coupled with different chromophores for easy identification of bands, with three reference proteins carrying enhanced intensity corresponding to a blue band at 20 kDa, green at 40 kDa, and red at 75 kDa, respectively, as separated on SDS-PAGE (Tris-Glycine buffer). The PM5000 ExcelBand™ 3-color Pre-Stained Protein Ladder Regular Range is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transferefficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.

Contents

Approximately $0.1^{\circ}0.4$ mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5 at 25°C), 2% SDS, 0.2 mM Dithiothreitol, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control


Under suggested conditions, the PM5000 ExcelBand™ 3-color Pre-Stained Protein Ladder Regular Range resolves 13 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.

Storage

4°C -20°C for 3 months for long term storage

ExcelBand™ 3-color Pre-Stained Protein Ladder, High Range (9-245 kDa)

PM5100 (250 μ l × 2)



Tris-Glycine

Description

The PM5100 ExcelBand™ 3-color Pre-Stained Protein Ladder High Range is a ready-to-use three-color protein standard with 14 pre-stained proteins covering a wide range of molecular weights from 10 to 245 kDa in Tris-Glycine Buffer (9 to 235 kDa in Bis-Tris (MOPS) buffer and 10 to 235 kDa in Bis-Tris (MES) buffer). Proteins are covalently coupled with different chromophores for easy identification of bands, with three reference proteins carrying enhanced intensity corresponding to a blue band at 20 kDa, green at 40 kDa, and red at 75 kDa, respectively, as separated on SDS-PAGE (Tris-Glycine buffer). The PM5100 3-color Pre-Stained Protein Ladder High Range is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.

Contents

Approximately $0.1^{\circ}0.4$ mg/ml of each protein in the buffer (20 mM Tris-phosphate (pH 7.5 at 25°C), 2% SDS, 0.2 mM Dithiothreitol, 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control

Under suggested conditions, the PM5100 ExcelBand™ 3-color Pre-Stained Protein Ladder High Range resolves 14 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.

Storage

4°C -20°C for 3 months

for long term storage

20°C










PM5200 (250 μl × 2)

Description



The PM5200 ExcelBand™ 3-color Pre-Stained Protein Ladder Broad Range is a ready-to-use three-color protein standard with 15 pre-stained proteins covering a wide range of molecular weights from 5 to 245 kDa in Tris-Glycine Buffer (3.5 to 235 kDa in Bis-Tris (MOPS) buffer and Bis-Tris (MES) buffer). Proteins are covalently coupled with different chromophores for easy identification of bands, with three reference proteins carrying enhanced intensity corresponding to a blue band at 20 kDa, green at 40 kDa, and red at 75 kDa, respectively, as separated on SDS-PAGE (Tris-Glycine buffer). The PM5200 3-color Pre-Stained Protein Ladder Broad Range is designed for monitoring protein separation during SDS-polyacrylamide gel electrophoresis, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the size of proteins.

Contents

Approximately $0.1^{\circ}0.4$ mg/ml of each protein in the buffer (20 mM Tris-phosphate, pH 7.5 at 25°C, 2% SDS, 0.2 mM Dithiothreit-ol 3.6 M Urea, and 15% (v/v) Glycerol).

Quality Control

Under suggested conditions, the PM5200 ExcelBand™ 3-color Pre-Stained Protein Ladder Broad Range resolves 15 major bands in SDS-PAGE (Tris-Glycine, MOPS, and MES buffer) and after Western blotting to a nitrocellulose membrane.

Storage

4°C -20°C

for 3 months for long term storage





YesBlot™ Western Marker I









WM1000 (250 μl)

Description

YesBlot™ Western Marker I is a ready-to-use mixture with ten IgG-binding proteins covering a wide range of molecular weights from 15 to 200 kDa in Tris- Glycine buffer. YesBlot™ Western Marker I performs dual functions. First, it contains 4 pre-stained proteins (10, 25, 45 and 70 kDa) for monitoring protein separation during SDS-PAGE, verification of Western transfer efficiency on membranes (nitrocellulose, PVDF, or nylon) and for approximating the protein size. Second, ten IgG-binding proteins can be immuno- detected on film or by CCD imaging.

YesBlot™ Western Marker I is compatible for chemiluminescent, fluorescent, chromogenic or other detection systems. In addition, YesBlot™ Western Marker I has two reference bands with enhanced intensity (at 30 kDa and 80 kDa).

The marker is supplied in gel loading buffer and is ready to use. Do NOT heat, dilute, or add reducing agents before loading.

Contents


The YesBlot™ Western Marker I contains recombinant IgG binding proteins, prestained recombinant proteins, Glycerol, and SDS.

Quality Control

Under suggested conditions, the YesBlot™ Western Marker I resolves 4 pre-stained bands on the membrane and 10 bands after primary and secondary antibodies binding followed by chemiluminescent detection.

Storage

4°C for 3 months
-20°C for long term storage





FluoroStain™ Protein Fluorescent Staining Dye (Red, 1000X)









PS1000 (1 ml × 1) PS1001 (1 ml × 5)

Description

FluoroStain™ Protein Fluorescent Staining (PS1000/PS1001) is designed to substitute the common Coomassie Blue protein staining method, offering greater sensitivity and ease of operation. Unlike Coomassie Blue stain, the FluoroStain™ Protein Fluorescent Staining Dye binds to protein with high specificity, making destaining process an option rather than a requirement. With further reduction of background signals via destaining process, the FluoroStain™ Protein Fluorescent Staining Dye is capable of achieving detection level parallel to silver stain without specialized imaging equipment (Fig. 1), making it one of the most sensitive dyes available. In addition to its remarkable sensitivity, the FluoroStain™ Protein Fluorescent Staining Dye brings a more reliable and safer user experience, since the stained gel can be visualized with blue-light illumination, users avoid the risk of skin/ eye damage caused by UV light. For best result, we suggest using the B-BOX™ Blue Light LED epi-illuminator to visualize and analyze the gel stained with FluoroStain™ Protein Fluorescent Staining Dye.

The FluoroStain™ Protein Fluorescent Staining Dye is compatible to the analysis of mass spectra, i.e. LC-MS/MS, MALDI-TOF, etc. (Fig. 2). The FluoroStain™ Protein Fluorescent Staining Dye is also for a less toxic (Fig. 3) and more environmentally-friendly procedure for protein staining, because it's designed to be used in a aqueous solution of ethanol and phosphoric acid for staining, avoiding the use of conventional methanol / acetic acid solution which is much more harmful and stimulating.

Spectral Characteristics

When it is bound with bovine serum albumin (BSA), the fluorescent emission of FluoroStain™ Protein Fluorescent Staining Dye can be excited by UV and blue light sources, with excitation peaks around 369 and 517 nm and emission at 605 nm (Fig. 4). In absence of BSA, FluoroStain™ Protein Fluorescent Staining Dye shows ignorable fluorescence as compared with protein-bound form, therefore giving a clear background for photographic analysis. These spectral characteristics made this fluorescent dye compatible with a wide variety of gel reading facilities, including UV/ blue light epi- and transilluminator, argon laser and mercury-arc lamp excitation gel scanners.




Fig.1. Remarkable sensitivity of the FluoroStain™ Protein Fluorescent Staining Dye (PS1000/PS1001) in comparison with Silver Stain.




Fig. 2. Comparison of MALDI-TOF mass spectra of bovine serum albumin (BSA) stained with FluoroStain™ Protein Fluorescent Staining Dye (A) and with Coomassie Blue (B). BSA proteins are seperated on an SDS-PAGE, stained with fluorescent dye or conventional Coomassive Blue, followed by trypsin digestion in gel, and then analyzed by MALDI-TOF.




Fig. 3. FluoroStain $^{\text{TM}}$ Protein Fluorescent Staining Dye shows no mutagenic activity in the Ames test.




Fig. 4. The Excitation and Emission Spectra of FluoroStain™ Protein Fluorescent Staining Dye (PS1000/PS1001) bound with BSA protein.

Working Reagent Preparation 1:1000 dilution in 40% ethanol and 2% H₃PO₄

Storage
Protected from light
-20°C for 24 months








Deoxynucleotide(dNTP)

dNTP Single Solution



Cat. No. **Product Description**

| CD1010 | Deoxynucleotide (dNTP) Mix, 10 mM each (40 mM total), 200 μl |
|--------|--|
| CD1011 | Deoxynucleotide (dNTP) Mix, 10 mM each (40 mM total), 200 μl x 5 |
| CD1020 | Deoxynucleotide (dNTP) Mix, 25 mM each (100 mM total), 500 μl |
| CD1021 | Deoxynucleotide (dNTP) Mix, 25 mM each (100 mM total), 500 μ l x 6 |
| CD3000 | dATP Solution - Sodium Salt (100 mM), 25 ml |
| CD3001 | dATP Solution - Sodium Salt (100 mM), 100 ml |
| CD4000 | dTTP Solution - Sodium Salt (100 mM), 25 ml |
| CD4001 | dTTP Solution - Sodium Salt (100 mM), 100 ml |
| CD5000 | dCTP Solution - Sodium Salt (100 mM), 25 ml |
| CD5001 | dCTP Solution - Sodium Salt (100 mM), 100 ml |
| CD6000 | dGTP Solution - Sodium Salt (100 mM), 25 ml |
| CD6001 | dGTP Solution - Sodium Salt (100 mM), 100 ml |
| CD7000 | dUTP Solution - Sodium Salt (100 mM), 25 ml |
| CD7001 | dUTP Solution - Sodium Salt (100 mM), 100 ml |







SMOChem™ dNTP Single Solution

Description

Highly quality dNTPs are a vital requirement for successful PCR, as the presence of contaminating impurities will result in a decrease in amplification sensitivity and yield of PCR product. SMOChem™ dNTP Single Solutions are ready-to-use dNTP aqueous solutions include 100 mM solutions of dATP, dCTP, dGTP, dTTP and dUTP respectively. SMOChem™ dNTPs undergo highly stringent purification steps to offer greater than 99% purity and are tested for the absence of DNase, RNase, Protease, and Nicking activity. dNTPs are suitable for application in PCR, sequencing, fill-in reactions,cDNA synthesis, nick translation, and TdT-tailing reactions. SMOChem™ dNTP Single Solution supplied as sodium salt in purified water (pH 8.5).


Features


- Greater than 99% purity determined by HPLC
- Ideal for use in PCR amplification and cDNA synthesis


- Long shelf-life
- Nuclease and ribonuclease free


Contents


| Component | Cat. No. | Volume |
|-------------------------------------|------------------|-----------------|
| dATP Solution- Sodium Salt (100 mM) | CD3000 CD3001 | 25 ml 100 ml |
| Component | Cat. No. | Volume |
| dTTP Solution- Sodium Salt (100 mM) | CD4000 CD4001 | 25 ml 100 ml |
| Component | Cat. No. | Volume |
| dCTP Solution- Sodium Salt (100 mM) | CD5000 CD5001 | 25 ml 100 ml |
| Component | Cat. No. | Volume |
| dGTP Solution- Sodium Salt (100 mM) | CD6000 CD6001 | 25 ml 100 ml |
| Component | Cat. No. | Volume |
| dUTP Solution- Sodium Salt (100 mM) | CD7000 CD7001 | 25 ml 100 ml |











SMOBIO's dNTPs show greater than 99% purity using HPLC analysis.

Storage

-20°C for 36 months





SMOChem™ dNTP Mix


Description


The SMOChem™ Deoxynucleotide (dNTP) Mix is an aqueous solution that contains an equimolar solution of ultrapure dATP, dCTP, dGTP and dTTP, each at a concentration of 10 mM or 25 mM at pH 8.5. The dNTP Mix is designed for many different molecular biology applications that involved in DNA synthesis or labeling, such as PCR, real-time PCR, DNA sequencing, reverse transcription, primer extension, and etc. The dNTP Mix is free of exo-deoxyribonuclease and endo-deoxyribonuclease as well as ribonuclease activity. The dNTP Mix offers the possibility to reduce the number of pipetting steps and the risk of reaction set up errors.


Features


- Greater than 99% purity determined by HPLC
- Ideal for use in PCR amplification and cDNA synthesis

- Long shelf-life
- Nuclease and ribonuclease free










SMOBIO'S dNTPs have highly performance for general PCR (A), long range PCR (B), RT-PCR (C), and real-time PCR (D).

Contents

| Component | Cat. No. | Volume |
|---|------------------|----------------------|
| Deoxynucleotide (dNTP) Mix, 10 mM each (40 mM total) | CD1010 CD1011 | 200 μl 200 μl × 5 |
| Component | Cat. No. | Volume |
| Deoxynucleotide (dNTP) Mix, 25 mM each (100 mM total) | CD1020 CD1021 | 500 μl 500 μl × 6 |

Storage

 -20°C for 36 months











DNA Amplification

High Fidelity PCR Standard PCR Extraction-free PCR Hot Start PCR Real-time PCR

SMOBIO DNA Polymerase Information

| | | High Fidelity PCR | | Standard PCR | Extraction free PCR | Hot Start PCR |
|------------------------------|------------|-------------------|----------------|---------------|---------------------|---------------|
| Catalog number | TF1000 | TF3000 | TK1000 | TP1000 | TP2000 | TP5000 |
| DNA polymerase | SMO-HiFi | G-HiFi | Klen-Taq | Taq | Blood direct | Hot Start II |
| | | | Propertio | es | | |
| Fidelity (comapred to Taq) | 70X | 70X | 4X | 1X | 1X | 1X |
| Amplification length | 12 kb | 40 kb | 10 kb | 8 kb | ≤ 2kb | 8 kb |
| Extenstion rate | 1 kb/ 10 s | 1 kb/ 7 s | 1 kb/ 20 s | 1 kb/ 20 s | 1 kb/ 20 s | 1 kb/ 20 s |
| Product end structure | blunt end | blunt end | 3'A/blunt end | 3′A | 3′A | 3'A |
| 3'→5' exonuclease activity | Yes | Yes | Yes | No | No | No |
| 5'→3' exonuclease activity | No | No | No | Yes | Yes | Yes |
| Jnits/ 50 μl reaction volume | 1U | 1U | 1.25U | 1.25~5U | 1.25~5U | 1.25~5U |
| Annealing temperature | Tm-5 | Tm-5 | Tm-5 | Tm-5 | Tm-5 | Tm-5 |
| | | | Applicatio | ons | | |
| Routine PCR | | ✓ | ✓ | * | ✓ | * |
| Colony PCR | | | | * | | * |
| High fidelity | * | * | ✓ | | | |
| High yield PCR | * | ✓ | ✓ | | | |
| High reaction rate | * | * | | | | |
| Long amplicon | ✓ | * | ✓ | | | |
| GC rich template | * | * | | | | |
| AT rich template | | | ✓ | ✓ | ✓ | |
| Multiplex PCR | | ✓ | | ✓ | ✓ | |
| Site-directed mutagenesis | | * | | | | |
| | | | Additional For | mats | | |
| Master Mix | | | | TP1100/TP1120 | | |
| Master Dye Mix | | | | TP1200 | TP2100 | |
| Fluorescent Master Mix | | | | TP1260 | | |

^{√ :} General use






^{★ :} Recommended choice

SMOBIO Q-PCR Master Mix Information

| D AVE | Product Name | ExcelTaq [™] 2X Q-PCR Master Mix | ExcelTaq™ 2X Q-PCR Master Mix | ExcelTaq™ 2X Fast Q-PCR Master | ExcelTaq™ 2X Fast Q-PCR Master | ExcelTaq™ 2X Q-PCR Master Mix |
|-------|-------------------------------------|---|--------------------------------|-----------------------------------|---------------------------------|--------------------------------|
| • | Cat. No. | (SYBR, no ROX) TQ1100/TQ1101 | (SYBR, ROX) TQ1110 | Mix (SYBR, no ROX) TQ1200/TQ1201 | Mix (SYBR, ROX) TQ1210/TQ1211 | (TaqMan, ROX) TQ2110 |
| | Detection chemistry | SYBR | SYBR | SYBR | SYBR | TaqMan |
| W | Blue contrast dye | ✓ | ✓ | ✓ | √ | х |
| | ROX reference dye | х | ✓ | х | ✓ | ✓ |
| 100 | Time required for enzyme activation | 10 min | 10 min | 2 min | 2 min | 10 min |
| | qPCR program | Standard | Standard | Fast and Standard | Fast and Standard | Standard |

How to choose SMOBIO Q-PCR Master Mix





SMO-HiFi™ DNA Polymerase









TF1000 (1 U/μl, 100 U)

Description

The SMO-HiFi™ DNA Polymerase is a new genetically modified, recombinant DNA polymerase with fidelity 70 times higher than *Taq* DNA polymerase during amplification, as well as very high elongation rate. Being highly thermostable, SMO-HiFi™ DNA Polymerase can remain viable even after being subjected to boiling for 2 minutes. The SMO-HiFi™ DNA Polymerase is also designed to operate in much lower Mg²+ concentration as compared to other DNA polymerase products.

Features

- 5'→3' DNA polymerase activity
- 3'→5' exonuclease (proofreading) activity
- High reaction rate (up to 1 kb/10 second)
- High fidelity, 70 times higher than Taq DNA polymerase
- Blunt end amplicons
- Thermo-stable: half-life is more than 10 hrs at 95°C

Contents

| Component | Volume |
|--|--|
| SMO-HiFi™ DNA Polymerase (1 U/µI) 10X HiFi™ Buffer 25mM MgSO ₄ dNTPs Mix (2 mM each) DMSO | 100 μl 600 μl 500 μl 600 μl 600 μl |
| DIVISO | 000 μι |

Storage Buffer

50 mM Tris-HCl (pH 8.0), 50 mM KCl, 0.1 mM EDTA, 1 mM DTT, stabilizers, and 50% (v/v) glycerol

Unit Definition

One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTPs into acid-insoluble material in 30 minutes at 74°C .

Storage




Fig. 1. SMO-HiFiTM DNA Polymerase's high processability enables reliable amplification of λDNA up to 12 kb in length (M: DM5100).




Fig. 2. SMO-HiFiTM DNA Polymerase can amplify PCR products from as little as 1 fg of template DNA (M: DM2000).

G-HiFi™ DNA Polymerase









TF3000 (1 U/μl, 100 U)

Description

The G-HiFi™ DNA Polymerase is a new genetically modified, recombinant DNA polymerase suitable for GC-rich templates that are difficult to amplify. The fidelity of G-HiFi™ DNA Polymerase is 70 times higher than that of Taq DNA polymerase. The high extension rate of G-HiFi™ DNA Polymerase is achieved by blending the DNA polymerase with an elongation enhancer. The optimized 5X G-HiFi™ Buffer includes special ingredients that suppress non-specific amplification as well as plateau effect produced by conventional PCR. With the optimized 5X G-HiFi™ Buffer, G-HiFi™ DNA Polymerase is capable to amplify most templates, such as longer targets (up to 40 kb from lambda DNA) and that contain GC-rich sequences.

Contents

| Component | Volume |
|---------------------------------|---------|
| G-HiFi™ DNA Polymerase (1 U/μl) | 100 μΙ |
| 5X G-HiFi™ Buffer | 1200 μΙ |
| dNTPs Mix (2 mM each) | 600 μΙ |

Features

- $5' \rightarrow 3'$ DNA polymerase activity
- 3'→5' exonuclease (proofreading) activity
- Suitable for GC-rich templates
- High reaction rate: 7 seconds/kb
- High fidelity: 70 times higher than Tag polymerase
- Generates blunt end amplicons
- Vast elongation capability (up to 40 kb)
- Thermo-stable for more than 10 hrs at 95°C.

Unit Definition

One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTPs into acid-insoluble material in 30 minutes at $74^{\circ}C$.

Storage Buffer

50 mM Tris-HCl (pH 8.0), 50 mM KCl, 0.1 mM EDTA, 1 mM DTT, stabilizers, and 50% (v/v) glycerol

Storage




Fig. 1. G-HiFiTM DNA Polymerase's high processability enables reliable amplification of λ DNA up to 40 kb in length (M: DM5100).




Fig. 2. G-HiFi[™] DNA Polymerase performs higher sensitivity for high GC content templates (GC: 71%) compare to high fidelity DNA Polymerase from Brand A (M: DM2000).

ExcelTaq™ Klen-Taq DNA Polymerase









TK1000 (5 U/μl, 500 U)

Description

The ExcelTaq $^{\text{TM}}$ Klen-Taq DNA Polymerase is a specially blended enzyme mix containing KlenTaq-1 DNA polymerase (a 5'-exo-minus, N-terminal deletion of Taq DNA polymerase) and a small amount of a proofreading DNA polymerase. This unique blending helps to improve the fidelity, yield and processivity of the resultant PCR process. Klen-Taq is also highly robust, showing high tolerance of varying concentrations of Mg^{2^+} ; it is highly thermostable and has four times the fidelity compared to Taq DNA polymerase. The ExcelTaq $^{\text{TM}}$ Klen-Taq DNA Polymerase is ideal for DNA amplifications 0.5-5 kb in length on genomic DNA, and up to 10 kb on less complex templates.

Features

- 5'→3' DNA polymerase activity
- 3'→5' exonuclease activity (proofreading)
- Thermo-stable: up to 98°C during PCR denaturing step
- 4X fidelity compared to Taq DNA polymerase
- Robust PCR performance, resistant to variance in PCR conditions

Contents

| Component | Volume |
|--|--------|
| ExcelTaq™ Klen-Taq DNA Polymerase (5 U/μl) | 100 μl |
| 10X Klen Buffer | 1.2 ml |

ExcelTag™ Klen-Tag Polymerase mixture

DNA polymerase, 5 units/μl Proofreading DNA polymerase Trace

Storage Buffer

40 mM Tris-HCl (pH 7.5), 50 mM KCl, 25 mM (NH $_4$) $_2$ SO $_4$, 0.1 mM EDTA, 5.0 mM 2-mercaptoethanol, stabilizer, 50% (v/v) glycerol

10X Klen Buffer

400 mM Tricine-KOH (pH 9.2), 150 mM KOAc, 35 mM Mg (OAc),, 750 μ g/ml BSA

Unit Definition

One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into an acid-insoluble material in 30 minutes at 74°C .

Storage




Fig. 1. ExcelTaq $^{\text{TM}}$ Klen-Taq DNA Polymerase can amplify PCR products from λ DNA up to 12 kb (M: DM5100).




Fig. 2. ExcelTaq™ Klen-Taq DNA Polymerase can amplify PCR products from as little as 1 pg of template DNA (M: DM2100).

ExcelTaq™ Taq DNA Polymerase









TP1000 (5 U/μl, 500 U)

Description

The ExcelTaq $^{\text{TM}}$ Taq DNA Polymerase is a recombinant thermo-stable DNA polymerase expressed and purified from an $E.\ coli$ strain carrying the cloned gene. With high DNA synthesis rate and thermo-stability, ExcelTaq $^{\text{TM}}$ DNA Polymerase is suitable for general and specialized PCR applications.

Features

- 5'→3' DNA polymerase activity
- 5'→3' exonuclease activity
- No detectable 3'→5' exonuclease (proofreading) activity
- Generates PCR products with 3'-dA overhangs
- Thermo-stable: half-life is more than 40 min at 95°C




Fig. 1. ExcelTaq $^{\text{m}}$ Taq DNA Polymerase can amplify PCR products from λ DNA up to 15 kb (M: DM3100).

Contents

| Component | Volume |
|--|----------|
| ExcelTaq™ <i>Taq</i> DNA Polymerase (5 U/μl) | 100 μl |
| 10X <i>Taq</i> Buffer | 1 ml x 2 |

Storage Buffer

20 mM Tris-HCl (pH 8.0), 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, Stabilizer, 50% (v/v) glycerol

10X Tag Buffer

200 mM Tris-HCl (pH 8.8), 100 mM KCl, 100 mM (NH $_4$) $_2$ SO $_4$, 20 mM MgCl,, 1% Triton X-100

Unit Definition

One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into an acid-insoluble material in 30 minutes at 74°C .

Storage

-20°C

for 24 months




Fig. 2. ExcelTaq $^{\text{m}}$ Taq DNA Polymerase can amplify PCR products from as little as 1 fg of template DNA (M: DM3100).

ExcelTaq™ PCR Master Mix









TP1100 (200 RXN) TP1120 (100 RXN)

Description

The ExcelTag™ PCR Master Mix is a ready-to-use mixture for amplifying targeted DNA fragments. It is designed to serve as ready-to-use master mix for virtually all PCR applications. The mixture contains all essential ingredients for PCR with the exception of template and primers. This not only saves valuable time in the laboratory, but also reduces the number of pipetting and reagent handling errors. The PCR Master Mix is supplied as a 5X/ 2X concentrated master mix, that is a mixture of recombinant Tag DNA polymerase, reaction buffer, MgCl₂ (TP1120 contains MgSO₄), dNTPs, and enzyme stabilizer.

Features

- 5'→3' DNA polymerase activity
- No detectable 3'→5' exonuclease (proofreading)
- Generates PCR products with 3'-dA overhangs
- High throughput PCR
- High Yield PCR
- High reproducibility, less pipetting errors

Contents

| TP1100 Component | Volume |
|-------------------------------------|-------------|
| ExcelTaq™ 5X PCR Master Mix | 1 ml x 2 |
| 6X DNA Loading Dye (Blue) | 1 ml x 2 |
| TP1120 Component | Volume |
| ExcelTaq™ 2X PCR Master Mix (MgSO₄) | 1.25 ml x 2 |
| 6X DNA Loading Dye (Blue) | 1 ml |

Storage

| 4°C | for 6 months |
|-------|---------------|
| -20°C | for 24 months |




Fig. 1. The ExcelTaq™ 5X PCR Master Mix can reliably amplify λDNA up to 8 kb in length (M: DM3100).





ExcelTaq™ 5x PCR Master Dye Mix









TP1200 (200 RXN)

Description

The ExcelTag[™] 5x PCR Master Dye Mix is a ready-to-use mixture for amplifying targeted DNA fragments. It is designed to serve as ready-to-use master mix for virtually all PCR applications. The mixture contains all essential ingredients for PCR with the exception of template and primers. This not only saves valuable time in the laboratory, but also reduces the number of pipetting and reagent handling errors. The PCR Master Dye Mix is supplied as a 5X concentrated ready-to-use mix, that is a mixture of recombinant Taq DNA Polymerase, reaction buffer, MgCl2, dNTPs, enzyme stabilizer and PCR friendly loading dye solution containing a tracking dye (Bromophenol blue) enabling efficient amplification of template in PCR and allowing the user to prepare a PCR reagent-loading dye master mix conveniently.

Features

- 5'→3' DNA polymerase activity
- No detectable 3'→5' exonuclease (proofreading) activity
- Generates PCR products with 3'-dA overhangs
- High throughput PCR
- High Yield PCR
- High reproducibility, less pipetting errors
- Load directly into electrophoresis

Contents

| Component | Volume | |
|---------------------------------|----------|--|
| ExcelTaq™ 5X PCR Master Dye Mix | 1 ml x 2 | |

Storage

4°C for 6 months -20°C for 24 months




Fig. 1. The ExcelTaq $^{\text{TM}}$ 5X PCR Master Dye Mix can reliably amplify λ DNA up to 8 kb in length (M: DM3100).











1 ml x 2

TP1260 (200 RXN)

Description

The ExcelTaq™ 5X Fluorescent PCR Master Mix is a ready-to-use mixture for amplifying targeted DNA fragments. It is designed to serve as ready-to-use master mix for virtually all PCR applications. The ExcelTaq™ 5X Fluorescent PCR Master Mix is supplied as a 5X concentrated ready-to-use mixture containing all the essential ingredients for PCR with the exception of template and primers. In addition, the mixture contains a tracking dye (Bromophenol blue), and a safer fluorescent DNA staining dye, which enables the user to track the electrophoresis process in real time as well as eliminating the need for staining process. The resultant PCR reaction mixture is sufficiently dense enough to be loaded directly into 1X TAE or 1X TBE buffer for electrophoresis.

Features

- 5'→3' DNA polymerase activity
- No detectable 3'→5' exonuclease (proofreading) activity
- Generates PCR products with 3'-dA overhangs
- High throughput PCR
- High Yield PCR
- High reproducibility, less pipetting errors
- Load directly into electrophoresis
- DNA bands can be visualized directly under UV or 470 nm blue light illumination

Contents


| Component | Volume |
|-----------|--------|
| | |

ExcelTag™ 5X Fluorescent PCR Master Mix

Storage

Protected from light and avoid multiple freeze/thaw cycles
4°C for 6 months


-20°C for 24 months



Viewed with B-Box™

Viewed with UV light

Fig. 1. λDNA was amplified with ExcelTaq $^{\text{TM}}$ 5X Fluorescent PCR Master Mix. The λDNA was amplified with specific primers for amplifying different ranges. The pictures were captured under B-BOX $^{\text{TM}}$ blue light (left). While using UV light (right), the amplicons can be seen in the right picture (M: DM 3160).



Viewed with B-Box™

Viewed with UV light

Fig. 2. Colony PCR with ExcelTaq™ 5X Fluorescent PCR Master Dye Mix. The JM109 or modified JM109 was used for amplifying the genomic galactosidase Z gene. The pictures were captured under B-BOX™ blue light (left). While using UV light (right), the amplicons can be seen in the right picture (M: DM 3160).

ExcelTag™ Blood Direct DNA Polymerase









TP2000 (5 U/µl, 500 U)

Description

The ExcelTaq™ Blood Direct DNA Polymerase is designed for amplifying targeted DNA directly from whole blood, eliminating the need for a lengthy DNA isolation process. The ExcelTaq™ Blood Direct DNA Polymerase is highly tolerant in the presence of PCR interfering/inhibiting substances in blood, such as IgG, hemoglobin, and lactoferrin. The ExcelTaq™ Blood Direct DNA Polymerase is compatible with most anticoagulants, such as citrate, EDTA, and heparin (Fig. 1). The ExcelTaq™ Blood Direct DNA Polymerase includes a pair of positive control primers (CCR5) that are compatible with primate blood samples.

Features

- 5'→3' DNA polymerase activity
- No detectable 3'→5' exonuclease (proofreading) activity
- Generates PCR products with 3'-dA overhangs
- Perform PCR directly from blood samples
- Compatible with most anticoagulants




Fig. 1. The ExcelTaq™ Blood Direct DNA Polymerase amplified 200 bp from separately treated blood samples using CCR5 specific primers. M: DM2100 marker, lane 1: fresh blood, lane 2: blood + citrate, lane 3: blood + EDTA, lane 4: blood + EDTA/ NaF, lane 5: blood + Heparin, lane 6: 1 mm² of dry blood on filter paper.

Contents

| Component | Volume | |
|--|--------------------|--|
| ExcelTaq™ Blood Direct DNA Polymerase (5 U/μl) 5X Blood Direct Buffer | 100 μl 1 ml x 4 | |
| Positive Control Primers (10 μM, each) | 50 μl | |

Storage Buffer

20 mM Tris-HCl (pH 8.0), 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, stabilizer, 50% (v/v) glycerol

Storage



ExcelTaq™ 5X Blood Direct PCR Master Mix Kit









TP2100 (200 RXN)

Description

The ExcelTaq™ 5X Blood Direct PCR Master Mix Kit is designed for amplifying targeted DNA directly from whole blood, eliminating the need for a lengthy DNA isolation process. The PCR master mix kits contains all the essential components for a PCR reaction and a PCR friendly loading and tracking dye (Orange G) allowing the user to easily prepare a PCR reagent and directly loading PCR product into agarose gel for electrophoresis.

The ExcelTaq™ 5X Blood Direct PCR Master Mix Kit is capable of tolerating the presence of PCR interfering/inhibiting substances in blood and is ideal for high-throughput screening of blood samples for high reproducibility. The PCR master mix kit includes a pairs of positive control primers (CCR5) that are compatible with primate blood samples.

Features

- 5'→3' DNA polymerase activity
- No detectable 3'→5' exonuclease (proofreading) activity
- Generates PCR products with 3'-dA overhangs
- High throughput PCR
- Execute PCR directly from blood samples
- High reproducibility, less pipetting errors
- Compatible with most anticoagulants

Contents

| Component | Volume | |
|--|----------|--|
| ExcelTaq™ 5X Blood Direct PCR Master Mix | 1 ml x 2 | |
| Positive Control Primers (10 μM, each) | 25 µl | |

Storage

4°C for 6 months -20°C for 24 months Caution: Avoid multiple freeze/thaw cycles



Fig. 1. The ExcelTaq™ 5X Blood Direct PCR Master Mix Kit amplified 200 bp from differently treated blood samples using CCR5 specific primers. M: DM2100 marker, Lane 1: fresh blood, Lane 2: blood + citrate, Lane 3: blood + EDTA, Lane 4: blood + EDTA/ NaF, Lane 5: blood + Heparin, Lane 6: 1mm² of dry blood on filter paper.











TP5000 (5 U/µl, 500 U)

Description

The ExcelTaq™ Hot Start II DNA Polymerase is a mixture of an aptamer-based inhibitor and a recombinant thermo-stable *Taq* DNA polymerase designed for preventing or minimizing non-specific DNA amplification in PCR reaction. The inactivation of polymerase is achieved by a reversible binding of the aptamer to the polymerase at temperatures below 45°C. The aptamer inhibitor releases polymerase during normal PCR cycling. The aptamer-based inhibition omits the time-consuming initial activation step required by chemically modified or antibody-based hot start polymerases.

The high specificity and sensitivity of ExcelTaq™ Hot Start II DNA Polymerase allows sensitive detection from limited amount of DNA templates, such as 1 pg of cDNA or 1 fg of plasmid DNA. With a high DNA synthesis rate and high thermo-stability, the ExcelTaq™ Hot Start II DNA Polymerase allows reactions to be set up at room temperature and is suitable for common and specialized PCR applications.

Features

- Reversible enzyme inactivation
- •Omits extra enzyme activation step
- Convenient for room temperature PCR set-up
- High yield and specificity of target amplicons
- Wide range of amplicon length (up to 10 kb)
- High sensitivity (as low as 1 fg of plasmid)
- Aptamer-based hot start PCR

Contents

| Component | Volume | |
|--------------------------------------|----------|--|
| Hot Start II DNA Polymerase (5 U/μΙ) | 100 μΙ | |
| 10X HS Buffer | 1 ml x 2 | |

Storage

-20°C for 24 months

Storage Buffer

50 mM Tris-HCl (pH 8.0), 50 mM KCl, 0.1 mM EDTA, 1 mM DTT, stabilizer, 50% (v/v) glycerol

10X HS Buffer

200 mM Tris-HCl (pH 8.8 at 25°C), 100 mM KCl, 100 mM (NH4)₂SO₄, 20 mM MgCl₂, 1% Triton X-100

Unit Definition

One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid-insoluble material in 30 minutes at 74°C .




Fig. 1. ExcelTaq™ Hot Start II DNA Polymerase shows high specificity on amplifying target DNA. The optimal annealing temperature of GAPDH primer set is 58°C. Improper annealing temperature set at 52°C may force primer-dimer formation. ExcelTaq™ Hot Start II DNA Polymerase eliminated primer-dimer and increased amounts of desired product at improper annealing temperature of 52°C.




Fig. 2. ExcelTaq $^{\text{TM}}$ Hot Start II DNA Polymerase shows high sensitivity to amplify from low amount of templates. Each set of PCR reactions contained either 1 pg, 10 pg, or 1 ng of HeLa cell cDNA as templates. ExcelTaq $^{\text{TM}}$ Hot Start II DNA Polymerase successfully amplified targets from lower amount of templates, in comparison with hot-start DNA polymerases from other suppliers and general Taq DNA polymerase

ExcelTaq™ 2X Q-PCR Master Mix (SYBR)







TQ1100 (200 RXN)(SYBR, no ROX) TQ1101(500 RXN)(SYBR, no ROX) TQ1110 (200 RXN)(SYBR, ROX)

Description

The ExcelTag™ 2X Q-PCR Master Mix (SYBR) is a ready-to-use reagent with all the essential components for quantitative real-time PCR (qPCR) except primers and template. The master mix features high sensitivity and signal intensity(Fig. 1-2) as well as low background and better compatibility with cDNA templates derived directly from reverse transcription reaction mixture.(Fig. 3) The ExcelTaq™ 2X Q-PCR Master Mix (SYBR) contains hot-start Tag polymerase in an optimized buffer with dsDNA specific SYBR green fluorescent dye. This master mix allows for sensitive, precise amplification, real-time tracking of the amplification process, and simultaneous quantification for targeted DNA molecules. With inert smart blue contrast dye, the ExcelTag™ 2X Q-PCR Master Mix (SYBR) is ready-to-use and greatly reduces pipetting errors, while largely improving the reproducibility of the process. The TQ1110 ExcelTag™ 2X Q-PCR Master Mix (SYBR, ROX) includes ROX reference dye for normalization of each qPCR assay.

Features

- High sensitivity and signal intensity
- Better compatibility for reverse transcription
- Smart blue contrast dye as a visual aid for reaction setup
- Low background
- With ROX reference dye (TQ1110)




Fig. 1. The amplification plot of real-time PCR with cDNA templates ranging from 25 ng to 1.49 fg in quantity, analyzed by using TQ1100 ExcelTaq™ 2X Q-PCR Master Mix (SYBR, no ROX) for qPCR amplification.

Contents

| TQ1100 Component | Volume |
|--|----------|
| ExcelTaq™ 2X Q-PCR Master Mix (SYBR, no ROX) | 1 ml x 2 |
| TQ1101 Component | Volume |
| ExcelTaq™ 2X Q-PCR Master Mix (SYBR, no ROX) | 1 ml x 5 |
| TQ1110 Component | Volume |
| ExcelTaq™ 2X Q-PCR Master Mix (SYBR, ROX) | 1 ml x 2 |

Storage

Aliquot to avoid multiple freeze-thaw cycles
Protect from light




Fig. 2. The amplification plot of real-time PCR with cDNA templates ranging from 1.49 fg to 25 ng in quantity, analyzed by using TQ1110 ExcelTaq™ 2X Q-PCR Master Mix (SYBR, ROX) for qPCR amplification.




Fig. 3. SMOBIO's TQ1100 ExcelTaq $^{\text{M}}$ 2X Q-PCR Master Mix (SYBR, no ROX) shows better compatibility with reverse-transcription reaction mixture as compared to similar products from A and B brands. Two μl of cDNA directly obtained from reverse-transcription reaction mixture were used in a 20 μl qPCR reaction for the compatibility test.



ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR)







TQ1200 (200 RXN)(SYBR, no ROX) TQ1201(500 RXN)(SYBR, no ROX) TQ1210 (200 RXN)(SYBR, ROX) TQ1211 (500 RXN)(SYBR, ROX)

Description

The ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR) is a ready-to-use reagent with all the essential components for quantitative real-time PCR (qPCR) except primers and templates. The master mix features high sensitivity and signal intensity as well as low background and better compatibility with fast PCR programs.

The ExcelTaq $^{\text{TM}}$ 2X Fast Q-PCR Master Mix (SYBR) contains hot-start Taq polymerase in an optimized buffer with dsDNA specific SYBR green fluorescent dye. This master mix allows sensitive, precise amplification, real-time tracking of the amplification process, and simultaneous quantification for targeted DNA molecules.

With inert smart blue contrast dye, the ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR) is ready-to-use and greatly reduces pipetting errors, while largely improving the reproducibility of the process. TQ1210 ExcelTaq™ 2X Fast Q-PCR Master(SYBR, ROX) includes ROX reference dye for normalization of each qPCR assay

Features

- · Fast hot start
- High stability
- High sensitivity and signal intensity
- Compatible with fast PCR program
- Smart blue contrast dye as a visual aid for reaction setup
- With ROX reference dye (TQ1210)




Fig. 1. The overlapped amplification curves from hot start duration of 2, 5, and 10 minutes display that TQ1200 ExcelTaq $^{\text{TM}}$ 2X Fast Q-PCR Master Mix (SYBR, no ROX) preforms successfully in short duration of initial activation (2 min).

ExcelTaq Q-PCR Master Mix features blue dye to clearly recognize aliquoted reaction mixes in plates to minimize aliquoting errors.

Contents

| TQ1200 Component | Volume |
|---|----------|
| ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR, no ROX) | 1 ml x 2 |
| TQ1201 Component | Volume |
| ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR, no ROX) | 1 ml x 5 |
| TQ1210 Component | Volume |
| ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR, ROX) | 1 ml x 2 |
| TQ1211 Component | Volume |
| ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR, ROX) | 1 ml x 5 |

Storage

Aliquot to avoid multiple freeze-thaw cycles Protect from light




Fig. 2. SMOBIO's TQ1200 ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR, no ROX) shows higher stability as compared to a similar products (Brand P and B).





ExcelTaq™ 2X Q-PCR Master Mix (TaqMan, ROX)









TQ2110 (200 RXN)

Description

The ExcelTaq[™] 2X Q-PCR Master Mix (TaqMan, ROX) is a ready-to-use reagent with all the essential components for quantitative real-time PCR (qPCR) except primers, TaqMan probes and templates. The master mix includes a 5′ to 3′ exonuclease activity to cleave TaqMan probes that hybridize to target sequences, releasing fluorophore during probe displacement. With TaqMan probes, the master mix features high specificity and high sensitivity (Fig. 1).

The ExcelTaq™ 2X Q-PCR Master Mix (TaqMan, ROX) contains hot-start *Taq* polymerase in an optimized buffer that allows for sensitive and precise amplification, real-time tracking of the amplification process, and simultaneous quantification for targeted DNA molecules. The master mix includes ROX reference dye for the normalization of each qPCR assay. The ExcelTaq™ 2X Q-PCR Master Mix (TaqMan, ROX) is ready-to-use and greatly reduces pipetting errors, while largely improving the reproducibility in the process.

Features

- · High sensitivity
- High specificity
- With ROX reference dye




Fig. 1. The amplification plot of real-time PCR with cDNA templates ranging from 76 fg to 20 ng in quantity, analyzed by using TQ2110 ExcelTaq $^{\text{TM}}$ 2X Q-PCR Master Mix (TaqMan, ROX) for qPCR amplification.

Contents

Component

Volume

ExcelTaq™ 2X Q-PCR Master Mix (TaqMan, ROX)

1 ml x 2

Storage

Aliquot to avoid multiple freeze-thaw cycles Protect from light

-20°C for 12 months




Fig. 2. The standard curve of TQ2110 ExcelTaq™ 2X Q-PCR Master Mix (TaqMan, ROX) was generated by using a 4-fold dilution of total cDNA ranging from 76 fg to 20 ng.







cDNA Synthesis and RT-PCR

cDNA Synthesis

One-Step RT-PCR

One-Step RT-qPCR

RNase Inhibitors







ExcelRT™ Reverse Transcriptase









Volume

Volume

RP1000 (200 U/µl, 20,000 U) RP1300 (100 RXN) RP1400 (100 RXN)

Description

The ExcelRT™ Reverse Transcriptase is a recombinant Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase - an RNA dependent DNA polymerase capable of generating first strand cDNA using an RNA template. It is designed to reduce RNase H activity and create better thermal stability. The $\mathsf{ExcelRT}^{\mathsf{TM}}$ Reverse Transcriptase is able to routinely synthesize first strand cDNA > 8 kb at 37~50°C.

Additional Kit Format

The ExcelRT™ Reverse Transcription Kits contain all components to synthesize high quality first strand cDNA. The kits contain ExcelRT™ Reverse Transcriptase, RNAok™ RNase Inhibitor, oligo (dT)₂₀ and random hexamers, which are used to synthesize cDNA from poly(A) tailed mRNA and total RNA, respectively. The RP1400 ExcelRT™ Reverse Transcription Kit II is supplied with Oligo (dT)/Random Primer Mix that is optimal for highly efficient synthesis of short chain cDNA suitable for real-time PCR.

Features

- · High yield
- Thermostable, up to 50°C, during first strand synthesis
- High processivity, generating cDNA up to 8 kb
- Reduced RNase H ribonuclease activity
- No detectable 3' → 5' exonucleolytic proofreading function
- Thermal stable for at least 4 weeks when stored at 4°C
- Suitable for real-time PCR
- Contains all components for reverse transcription (RP1300 and RP1400)
- Time saving for short chain cDNA synthesis (RP1400)




Fig. 1. ExcelRT™ Reverse Transcriptase can synthesize first strand cDNA using only 5 fg of total RNA in solution. The bands in the gel represent the results of GAPDH (230 bp) from two step RT-PCR reaction. (M: DM2100)

Contents

| RP1000 | | | |
|----------|---------|------------|----|
| ExcelRT™ | Reverse | Transcript | ta |

| Reverse Transcriptase (200 U/µI) 100 5X RT Buffer 1 | |
|---|-----|
| 0.1 M DTT 500 | m I |

RP1300

Component

ExcelRT™ Reverse Transcription Kit

| Reverse Transcriptase (200 U/μl) | 100 μΙ |
|----------------------------------|----------|
| RNase Inhibitor (20 U/μΙ) | 100 μΙ |
| 5X RT Buffer (DTT) | 500 μΙ |
| dNTPs (10 mM each) | 200 μΙ |
| Oligo (dT)20 (50 μM) | 100 μΙ |
| Random Hexamers (100 μM) | 100 μΙ |
| DEPC-Treated H ₂ O | 1 ml x 2 |

RP1400

ExcelRT™ Reverse Transcription Kit II Component

| RTase/RI Enzyme Mix | 100 μΙ |
|-------------------------------|----------|
| 5X RT Buffer (DTT/dNTPs) | 500 μl |
| Oligo (dT)/Random Primer Mix | 100 µl |
| DEPC-Treated H ₂ O | 1 ml x 2 |

Unit definition

One unit is defined as the amount of enzyme that will incorporate 1 nM of dTTP into acid-insoluble material in 10 minutes at 37°C using Poly (A) • oligo(dT)₂₅ as a template-primer.

Storage

-20°C for 24 months




Fig. 2. cDNA synthesized in 10 minutes is suitable for real-time PCR analysis. cDNA was synthesized in 10, 30, and 50 min from reverse transcription reaction. The amplification plot represents the results of 18S rRNA analyzed by using the ExcelTag™ 2X Q-PCR Master Mix (Cat. No. TQ1100) for real-time PCR amplification.



ExcelRT™ One-Step RT-PCR Kit










RP1100 (50 RXN)

Description

The ExcelRT™ One-Step RT-PCR Kit is designed for the reverse transcription and PCR amplification of a specific target RNA from either total RNA or mRNA. The ExcelRT™ One-Step RT-PCR Kit provides the user an alternative to the lengthy two step process (first strand generation and amplification) by using a single mixture, single tube, one step reaction. The ExcelRT™ One-Step RT-PCR Kit contains a 2X reaction premix consisting of an optimized buffer, dNTPs, Mg²⁺ and enzyme stabilizer, and a blend of recombinant reverse transcriptase and Taq DNA polymerase. The ExcelRT™ One-Step RT-PCR Kit allows the user to complete the RT-PCR process using a thermocycler in a single reaction setting and is ideal for target RNA amplification/ analysis capable of detecting even trace amounts of target RNA.

Features

- Generates dsDNA directly from RNA samples
- High throughput
- High reproducibility, less pipetting errors
- High sensitivity and yields
- Reduced RNase H ribonuclease activity






Fig. 1. ExcelRT™ One-Step RT- PCR Kit can synthesize PCR products with a minimal amount of total RNA in reaction solution. The bands in the gel represent the results of GAPDH (230 bp) from one-step RT-PCR reaction (M: DM2300)

Contents

| Component | Volume |
|--------------------|------------|
| 2X One-Step Buffer | 750 μl x 2 |
| Taq/RT Enzyme Mix | 50 μl |

Storage

-20°C for 24 months



ExcelRT™ One-Step RT-qPCR Kit (TagMan, ROX)









RQ2110 (200 RXN)

Description

The ExcelRT™ One-Step RT-qPCR kit (TaqMan, ROX) is designed for reverse transcription and quantitative real-time analysis of a specific target RNA by one-step reaction. The ExcelRT™ One-Step RT-gPCR kit (TagMan, ROX), consisting of One-Step RT Enzyme Mix and 2X One-Step Master Mix, is a convenient kit designed for highly efficient cDNA synthesis and highly specific real-time PCR in a single tube. The One-Step RT Enzyme Mix contains a thermostable ExcelRT™ Reverse Transcriptase and a RNAok™ RNase inhibitor. Consequently, One-Step RT Enzyme Mix can reverse transcribe RNA to cDNA at a wide temperature range from 42 to 60°C and be active against RNase A, RNase B and RNase C. By containing specialized hot-start Tag DNA polymerase, which greatly reduce primer-dimer formation and can be activated within 2 minutes, the 2X One-Step Master Mix features high specificity and is suitable for fast cycle program. This master mix includes ROX reference dye for normalization of each RT-qPCR assay.

Features

- High yield
- Reverse transcription at wide temperature range (42-60°C)
- High specificity
- Suitable for fast cycle program
- ROX reference dye





Fig. 1. ExcelRT™ One-Step RT-qPCR Kit can quantitatively analyze target RNA from a wide range of RNA template input. The amplification plot of one-step RT-qPCR with total RNA templates ranging from 1 pg to 1 ng in quantity, analyzed by using RQ2110 ExcelRT™ One-Step RT-qP-CR Kit (TagMan, ROX) for RT-qPCR amplification.

Contents

| Component | volume | |
|--------------------------------------|----------|--|
| One-Step RT Enzyme Mix | 400 μl | |
| 2X One-Step Master Mix (TagMan, ROX) | 1 ml x 2 | |

Storage

Aliquot to avoid multiple freeze-thaw cycles (stable within 30 freeze-thaw cycles) Protect from light -20°C for 12 months






Fig. 2. ExcelRT™ One-Step RT-qPCR Kit can quantitatively analyze target RNA at a wide temperature range (42-60°C). The amplification plot of one-step RT-qPCR with reverse transcription at temperature range from 42 to 60°C, analyzed by using RQ2110 ExcelRT™ One-Step RT-qPCR Kit (A) or kit from brand N (B) for RT-qPCR amplification. The overlapped amplification curves display that ExcelRT™ One-Step RT-qPCR Kit preforms successfully cDNA synthesis at wide temperature range.



ExcelRT™ One-Step RT-qPCR Kit (TaqMan, no ROX)

RQ2200 (200 RXN)

Description

The ExcelRT™ One-Step RT-qPCR kit (TaqMan, no ROX) is designed for reverse transcription and quantitative real-time analysis of a specific target RNA by one-step reaction. The ExcelRT™ One-Step RT-qPCR kit (TaqMan, no ROX), consisting of One-Step RT Enzyme Mix and 2× One-Step Master Mix, is a convenient kit designed for highly efficient cDNA synthesis and highly specific real-time PCR in a single tube. The One-Step RT Enzyme Mix contains a thermostable ExcelRT™ Reverse Transcriptase and a RNAok™ RNase inhibitor. Consequently, One-Step RT Enzyme Mix can reverse transcribe RNA to cDNA at a wide temperature range from 42 to 60°C and be active against RNase A, RNase B and RNase C. By containing specialized hot-start Tag DNA polymerase, which greatly reduce primer-dimer formation and can be activated within 2 minutes, the 2× One-Step Master Mix features high specificity and is suitable for fast cycle program.

Features


- High specificity
- With no ROX reference dye
- Suitable for fast program
- Reverse transcription at wide temperature range (42°C-60°C)

Contents

| Component | Volume | |
|---|----------|--|
| One-Step RT Enzyme Mix | 400 μl | |
| 2X One-Step Master Mix (TagMan, no ROX) | 1 ml x 2 | |

Storage

Aliquot to avoid multiple freeze-thaw cycles Protect from light -20°C for 12 months






Fig. 1. ExcelRT™ One-Step RT-qPCR Kit can quantitatively analyze target RNA from a wide range of RNA template input. The amplification plot of one-step RT-qPCR with total RNA templates ranging from 1 pg to 1 ng in quantity, analyzed by using RQ2200 ExcelRT™ One-Step RT-qP-CR Kit (TaqMan, no ROX) for RT-qPCR amplification.

Fig. 2. ExcelRT™ One-Step RT-qPCR Kit can quantitatively analyze target RNA at a wide temperature range (42-60°C). The overlapped amplification plot of one-step RT-qPCR with reverse transcription at temperature range from 42 to 60°C, analyzed by using RQ2200 ExcelRT™ One-Step RT-qPCR Kit display that ExcelRT™ One-Step RT-qPCR Kit performs successfully cDNA synthesis at wide temperature range.

RNAok™ RNase Inhibitor

RI1000 (20 U/µl, 2000 U x 1) RI1001 (20 U/µI, 2000 U x 5)

Description

RNAok™ RNase Inhibitor is a recombinant mammalian RNase inhibitor that is purified by affinity chromatography from E. coli. This protein inhibits pancreatic-type ribonucleases, RNase A, B, and C by binding strongly to RNases in a noncompetitive mode at a 1:1 ratio. RNAok™ RNase Inhibitor does not inhibit eukaryotic RNases T1, T2, U1, U2, CL3 as well as prokaryotic RNases I and H. RNAok™ RNase Inhibitor is compatible with RT-PCR enzymes such as AMV, M-MLV and ExcelRT™ Reverse Transcriptase or *Tag* DNA polymerase.




Fig. 1. RNAok™ RNase Inhibitor inhibits RNase A activity. Total HeLa RNA (1 µg) was incubated with 1U/ 1 µl of RNAok™ RNase Inhibitor and 0.4 ng RNase A for 30 min at 37°C. The result shows that RNAok™ RNase Inhibitor can protect RNA against 0.4 ng RNase A. M: DM3100, Lane 1: total HeLa RNA (1 µg), Lane 2: total HeLa RNA with RNase A (no inhibitor added), Lane 3: total HeLa RNA with RNase A and inhibitor from brand A, Lane 4-6: total HeLa RNA with RNase A and RNAok™ RNase Inhibi-

Application

- RT-PCR
- cDNA Synthesis
- in vitro translation
- in vitro transcription
- Separation and identification of specific ribonuclease activities

Usage Recommendation

Add RNAok™ RNase Inhibitor to transcription, translation, and cDNA synthesis reactions at a final concentration of 1 Unit/µl.

Storage Buffer

40 mM HEPES-KOH (pH 7.5), 100 mM KCl, 8 mM DTT, 0.1 mM EDTA, stabilizer and 50% (v/v) glycerol

Unit Definition

One unit is defined as the amount of RNAok™ RNase Inhibitor required to inhibit the activity of 5 ng of RNase A by 50%.

Storage

-20°C for 24 months





COVID-19

Real-Time RT-PCR Kit

For Research Use Only

Cat no. IP2000




INTENDED USE


For research purpose, IdPath™ COVID-19 Real-Time RT-PCR Kit is a real-time RT-PCR test intended for the detection of SARS-CoV-2 virus RNA which might be extracted from the respiratory tract specimens. The Kit provides reagents for multiplex real-time RT-PCR to detect SARS-CoV-2 by one-step reaction, specifically targeting the E (Envelope), RdRP (RNA-dependent RNA polymerase) and N (Nucleocapsid protein) gene for SARS-CoV-2 virus.


The Kit contains the RT enzyme mix and qPCR master mix for reverse transcription and real-time PCR of virus RNA. The COVID-19 Control (positive control) and ddWater (negative control) are used as indicators to avoid false negative/positive results across all experimental procedures. The Primers/probes Mix contains multiplex primers and TaqMan probes specific to the N, E/RdRP genes of SARS-CoV-2 and RNase P gene of human, detected by FAM, VIC and ROX channels, respectively.

FEATURES

- **High Sensitivity**: 5×10² copies/ml (10 copies/rxn)
- High Inclusivity: >99% of currently available complete virus genomes for SARS-CoV-2 including Omicron variant
- **High Accuracy** : Clinical validation with 100% accuracy
- High Compatibility : Suitable for most laboratory qPCR machines
- Operation Control: Including internal control for quality control of total process
- Convenience : Multiplex (E/RdRP and N) detection by one-step reaction







UD: Undetectable

NTC: Negative control (ddWater)

PC: Positive control for the SARS-COV-2 E/RdRP, N, and human RNase P genes.

ORDERING INFORMATION

| Catalog No. | Description | Package |
|-------------|--|-----------|
| IP2000 | IdPath™ COVID-19 Real-Time RT-PCR Kit (Cat no. IP2000) | 100 tests |

Small Bio, Smart Tool

Illuminator

| Cat. No. | Product Description |
|----------|---|
| VE0100 | B-BOX™ Blue Light LED Epi-illuminator, DC 12 Volt, 0.72 Amp |
| VE0101 | Adapter, AC 100-240V, 50/60 Hz, 1.8 m Power Cord, 1/pk |
| VE0102 | Multi-angle Filter Plate for B-BOX™, 1/pk |
| VE0103 | Amber Goggles for B-BOX™, 1/pk |
| VE0104 | Phox™ Photobox, 1/pk |
| Cloning | |
| Cat. No. | Product Description |
| CV1100 | GetClone™ PCR Cloning Vector II, 20 RXN |
| CK1000 | Champion™ E. coli Transformation Kit, 200 RXN |

DNA Ladders and Dyes

DNA Ladders and Markers

| Cat. No. | Product Description |
|----------|---|
| DM1100 | ExcelBand™ 50 bp DNA Ladder, 500 μl |
| DM2100 | ExcelBand™ 100 bp DNA Ladder, 500 μl |
| DM2300 | ExcelBand™ 100 bp+3K DNA Ladder, 500 μl |
| DM3100 | ExcelBand™ 1 KB (0.25-10 kb) DNA Ladder, 500 μl |
| DM3200 | ExcelBand™ 1 KB Plus (0.1-10 kb) DNA Ladder, 500 μl |
| DM4100 | ExcelBand™ XL 25 kb DNA Ladder, Broad Range (up to 25 kb), 500 μl |
| DM5100 | ExcelBand™ Super Range DNA Ladder (50 bp-25 kb) , 500 μl |
| DM1200 | AccuBand™ 50 bp DNA Ladder II, 500 μl |
| DM2000 | AccuBand™ 100 bp DNA Marker II, 500 μl |
| DM2200 | AccuBand™ 100 bp DNA Ladder II, 500 μl |

DNA Fluorescent Ladders

DM2400

| Cat. No. | Product Description |
|----------|--|
| DM1160 | FluoroBand™ 50 bp Fluorescent DNA Ladder, 500 μl |
| DM2160 | FluoroBand™ 100 bp Fluorescent DNA Ladder, 500 μl |
| DM2360 | FluoroBand™ 100 bp+3K Fluorescent DNA Ladder, 500 μl |
| DM3160 | FluoroBand™ 1 KB (0.25-10 kb) Fluorescent DNA Ladder, 500 μl |
| DM3260 | FluoroBand™ 1 KB Plus (0.1-10 kb) Fluorescent DNA Ladder, 500 μl |
| DM4160 | FluoroBand™ XL 25 kb Fluorescent DNA Ladder, Broad Range (up to 25 kb), 500 µl |

AccuBand™ 100 bp+3K DNA Ladder II, 500 μl

DNA Loading Dyes

| Cat. No. | Product Description |
|----------|---|
| DL1000 | ExcelDye™ 6X DNA Loading Dye, Orange, 5 ml x 2 |
| DL2000 | ExcelDye™ 6X DNA Loading Dye, Green, 5 ml x 2 |
| DL3000 | ExcelDye™ 6X DNA Loading Dye, Blue, 5 ml x 2 |
| DL4000 | ExcelDye™ 6X DNA Loading Dye, Tri-color, 5 ml x 2 |

DNA Fluorescent Loading Dye

| Cat. No. | Product Description |
|----------|--|
| DL5000 | FluoroDye™ DNA Fluorescent Loading Dye (Green, 6X), 1 ml |
| DL5001 | FluoroDye™ DNA Fluorescent Loading Dye (Green, 6X), 1 ml x 5 |



DNA Fluorescent Staining Dyes

| Cat. No. | Product Description |
|----------|---|
| DS1000 | FluoroStain™ DNA Fluorescent Staining Dye (Green, 10,000X), 500 µl |
| DS1001 | FluoroStain $^{\text{\tiny TM}}$ DNA Fluorescent Staining Dye (Green, 10,000X), 500 μl x 5 |
| NS1000 | FluoroVue™ Nucleic Acid Gel Stain (10,000X), 500 μl |
| NS1001 | FluoroVue™ Nucleic Acid Gel Stain (10,000X), 500 µl x 5 |
| | |

Precast Gel

Bis-Tris series

| Cat. No. | Product Description |
|----------|---|
| QP2110 | Q-PAGE™ Bis-Tris Precast Gel (Mini, 12 wells, 8%), 10 gels |
| QP2310 | Q-PAGE™ Bis-Tris Precast Gel (Mini, 12 wells, 12%), 10 gels |
| QP2510 | Q-PAGE™ Bis-Tris Precast Gel (Mini, 12 wells, 4-12%), 10 gels |
| QP3110 | Q-PAGE™ Bis-Tris Precast Gel (Midi, 12 wells, 8%), 10 gels |
| QP3310 | Q-PAGE™ Bis-Tris Precast Gel (Midi, 12 wells, 12%), 10 gels |
| QP3510 | Q-PAGE™ Bis-Tris Precast Gel (Midi, 12 wells, 4-12%), 10 gels |
| QP2120 | Q-PAGE™ Bis-Tris Precast Gel (Mini, 15 wells, 8%), 10 gels |
| QP2320 | Q-PAGE™ Bis-Tris Precast Gel (Mini, 15 wells, 12%), 10 gels |
| QP2520 | Q-PAGE™ Bis-Tris Precast Gel (Mini, 15 wells, 4-12%), 10 gels |
| QP3120 | Q-PAGE™ Bis-Tris Precast Gel (Midi, 15 wells, 8%), 10 gels |
| QP3320 | Q-PAGE™ Bis-Tris Precast Gel (Midi, 15 wells, 12%), 10 gels |
| QP3520 | Q-PAGE™ Bis-Tris Precast Gel (Midi, 15 wells, 4-12%), 10 gels |

TGN series

| Cat. No. | Product Description |
|----------|--|
| QP4220 | Q-PAGE™ TGN Precast Gel (Mini, 15 wells, 10%), 10 gels |
| QP4510 | Q-PAGE™ TGN Precast Gel (Mini, 12 wells, 4-15%), 10 gels |
| QP5210 | Q-PAGE™ TGN Precast Gel (Midi, 12 wells, 10%), 10 gels |
| QP5510 | Q-PAGE™ TGN Precast Gel (Midi, 12 wells, 4-15%), 10 gels |
| QP4220 | Q-PAGE™ TGN Precast Gel (Mini, 15 wells, 10%), 10 gels |
| QP4520 | Q-PAGE™ TGN Precast Gel (Mini, 15 wells, 4-15%), 10 gels |
| QP5220 | Q-PAGE™ TGN Precast Gel (Midi, 15 wells, 10%), 10 gels |
| QP5520 | Q-PAGE™ TGN Precast Gel (Midi, 15 wells, 4-15%), 10 gels |
| | |

Running Buffer Powder

| Cat. No. | Product Description |
|----------|---|
| BF2000 | SMOChem™ 1X MOPS-SDS Running Buffer Powder, 5 packs |
| BF4000 | SMOChem™ 1X Tris-Glycine-SDS Running Buffer Powder, 5 packs |

Protein Markers and Stain

Protein Markers and Ladders

| Cat. No. | Product Description |
|----------|---|
| PM1500 | ExcelBand™ All Blue Regular Range Protein Marker, 250 μl x 2 |
| PM1600 | ExcelBand™ All Blue Regular Range Plus Protein Marker, 250 μl x 2 |
| PM1700 | ExcelBand™ All Blue Broad Range Protein Marker, 250 μl x 2 |
| PM2500 | ExcelBand™ 3-color Regular Range Protein Marker, 250 μl x 2 |
| PM2510 | ExcelBand™ Enhanced 3-color Regular Range Protein Marker, 250 µl x 2 |
| PM2600 | ExcelBand™ 3-color High Range Protein Marker, 250 μl x 2 |
| PM2610 | ExcelBand™ Enhanced 3-color High Range Protein Marker, 250 µl x 2 |
| PM2700 | ExcelBand™ 3-color Broad Range Protein Marker, 250 μl x 2 |
| PM2800 | ExcelBand™ 3-color Extra Range Protein Marker, 250 μl x 2 |
| PM5000 | ExcelBand $^{\text{\tiny TM}}$ 3-color Pre-stained Protein Ladder, Regular Range, 250 μl x 2 |
| PM5100 | ExcelBand™ 3-color Pre-stained Protein Ladder, High Range, 250 µl x 2 |
| PM5200 | ExcelBand™ 3-color Pre-stained Protein Ladder, Broad Range, 250 µl x 2 |

Western Marker

| Cat. No. | Product Description |
|----------------------------------|--|
| WM1000 | YesBlot™ Western Marker I, 250 μl |
| Protein Fluorescent Staining Dye | |
| Cat. No. | Product Description |
| PS1000 | FluoroStain™ Protein Fluorescent Staining Dye (Red, 1000X), 1 ml |
| PS1001 | FluoroStain™ Protein Fluorescent Staining Dye (Red, 1000X), 1 ml x 5 |

Nucleic acid

| Cat. No. | Product Description |
|----------|--|
| CD1010 | Deoxynucleotide (dNTP) Mix, 10 mM each (40 mM total), 200 μl |
| CD1011 | Deoxynucleotide (dNTP) Mix, 10 mM each (40 mM total), 200 μ l x 5 |
| CD1020 | Deoxynucleotide (dNTP) Mix, 25 mM each (100 mM total), 500 μ l |
| CD1021 | Deoxynucleotide (dNTP) Mix, 25 mM each (100 mM total), 500 μ l x 6 |
| CD3000 | dATP Solution - Sodium Salt (100 mM), 25 ml |
| CD3001 | dATP Solution - Sodium Salt (100 mM), 100 ml |
| CD4000 | dTTP Solution - Sodium Salt (100 mM), 25 ml |
| CD4001 | dTTP Solution - Sodium Salt (100 mM), 100 ml |
| CD5000 | dCTP Solution - Sodium Salt (100 mM), 25 ml |
| CD5001 | dCTP Solution - Sodium Salt (100 mM), 100 ml |
| CD6000 | dGTP Solution - Sodium Salt (100 mM), 25 ml |
| CD6001 | dGTP Solution - Sodium Salt (100 mM), 100 ml |
| CD7000 | dUTP Solution - Sodium Salt (100 mM), 25 ml |
| CD7001 | dUTP Solution - Sodium Salt (100 mM), 100 ml |



DNA Amplification

| Hi | σh | Fide | litv | PCR |
|----|----|------|------|-----|
| | | | | |

| Cat. No. | Product Description |
|----------|---|
| TF1000 | SMO-HiFi™ DNA Polymerase, 1 U/µl, 100 U |
| TF3000 | G-HiFi™ DNA Polymerase, 1 U/μl, 100 U |
| TK1000 | ExcelTaq™ Klen-Taq DNA Polymerase, 5 U/μl , 500 U |
| | |

Standard PCR

| Cat. No. | Product Description |
|----------|---|
| TP1000 | ExcelTaq™ DNA Polymerase, 5 U/μl, 500 U |
| TP1100 | ExcelTaq™ 5X PCR Master Mix, 200 RXN |
| TP1120 | ExcelTaq [™] 2X PCR Master Mix (MgSO ₄), 100 RXN |
| TP1200 | ExcelTaq [™] 5X PCR Master Dye Mix, 200 RXN |
| TP1260 | ExcelTaq™ 5X Fluorescent PCR Master Mix, 200 RXN |

Extraction-free PCR

| Cat. No. | Product Description |
|----------|---|
| | |
| TP2000 | ExcelTaq™ Blood Direct DNA Polymerase, 5 U/μl, 500 U |
| TP2100 | ExcelTaq™ 5X Blood Direct PCR Master Mix Kit, 200 RXN |

Hot Start PCR

| Cat. No. | Product Description |
|----------|--|
| TP5000 | ExcelTaq™ Hot Start II DNA Polymerase, 5 U/μl, 500 U |

Real-time PCR

| Cat. No. | Product Description |
|----------|--|
| TQ1100 | ExcelTaq™ 2X Q-PCR Master Mix, (SYBR, no ROX), 200 RXN |
| TQ1101 | ExcelTaq™ 2X Q-PCR Master Mix, (SYBR, no ROX), 500 RXN |
| TQ1110 | ExcelTaq™ 2X Q-PCR Master Mix, (SYBR, ROX), 200 RXN |
| TQ1200 | ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR, no ROX), 200 RXN |
| TQ1201 | ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR, no ROX), 500 RXN |
| TQ1210 | ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR, ROX), 200 RXN |
| TQ1211 | ExcelTaq™ 2X Fast Q-PCR Master Mix (SYBR, ROX), 500 RXN |
| TQ2110 | ExcelTaq™ 2X Q-PCR Master Mix, (TaqMan, ROX), 200 RXN |

cDNA Synthesis and RT-PCR

cDNA Synthesis

| Cat. No. | Product Description |
|----------|---|
| RP1000 | ExcelRT™ Reverse Transcriptase, 200 U/μl, 20000 U |
| RP1300 | ExcelRT™ Reverse Transcription Kit, 100 RXN |
| RP1400 | ExcelRT™ Reverse Transcription Kit II, 100 RXN |
| | |

One-Step RT-PCR

| Cat. No. | Product Description | |
|----------|--------------------------------------|--|
| | | |
| RP1100 | ExcelRT™ One-Step RT-PCR Kit, 50 RXN | |
| | | |

One-Step RT-qPCR

| Cat. No. | Product Description |
|----------|---|
| RQ2110 | ExcelRT™ One-Step RT-qPCR Kit (TaqMan, ROX), 200 RXN |
| RQ2200 | ExcelRT™ One-Step RT-qPCR Kit (TaqMan, no ROX), 200 RXN |

RNase Inhibitor

| Cat. No. | Product Description |
|----------|---|
| RI1000 | RNAok™ RNase Inhibitor, 20 U/μl, 2000 U |
| RI1001 | RNAok™ RNase Inhibitor, 20 U/µl, 2000 U x 5 |

Pathogen Detection

| Cat. No. | Product Description |
|----------|--|
| | |
| IP2000 | IdPath™ COVID-19 Real-Time RT-PCR Kit, 100 RXN |









www.smobio.com